

Onasemnogene abeparvovec (exceeding € 50 million turnover limit: Spinal muscular atrophy)

Resolution of: 4 November 2021 Entry into force on: 4 November 2021 BAnz AT 07 12 2021 B3 Valid until: unlimited

Therapeutic indication (according to the marketing authorisation of 18 May 2020):

Zolgensma is indicated for the treatment of:

- patients with 5q spinal muscular atrophy (SMA) with a biallelic mutation in the SMN1 gene and a clinical diagnosis of SMA Type 1, or
- patients with 5q SMA with a biallelic mutation in the SMN1 gene and up to 3 copies of the SMN2 gene

Therapeutic indication of the resolution (resolution of 4 November 2021):

- see therapeutic indication according to marketing authorisation
- **1.** Additional benefit of the medicinal product in relation to the appropriate comparator therapy
- a) Patients with 5q spinal muscular atrophy (5q SMA) type 1

Appropriate comparator therapy:

Nusinersen

Extent and probability of the additional benefit of onasemnogene abeparvovec compared to the appropriate comparator therapy:

An additional benefit is not proven

b) Patients with 5q SMA type 2 and up to 3 copies of the SMN2 gene

Appropriate comparator therapy:

Nusinersen

Extent and probability of the additional benefit of onasemnogene abeparvovec compared to the appropriate comparator therapy

An additional benefit is not proven

c) Patients with 5q SMA type 3 and up to 3 copies of the SMN2 gene

Appropriate comparator therapy:

Treatment according to the doctor's instructions with the selection of nusinersen or BSC

Extent and probability of the additional benefit of onasemnogene abeparvovec compared to the appropriate comparator therapy:

An additional benefit is not proven

d) Pre-symptomatic patients with 5q SMA and up to 3 copies of the SMN2 gene

Appropriate comparator therapy:

Nusinersen

Extent and probability of the additional benefit of onasemnogene abeparvovec compared to the appropriate comparator therapy:

An additional benefit is not proven

Study results according to endpoints:

a) Patients with 5q SMA type 1

There are no assessable data.

Summary of results for relevant clinical endpoints

| Endpoint category | Direction of effect/ | Summary |
|--|----------------------|-------------------------------|
| | Risk of bias | |
| Mortality | n.a. | There are no assessable data. |
| Morbidity | n.a. | There are no assessable data. |
| Health-related quality | Ø | No data available. |
| of life | | |
| Side effects | n.a. | There are no assessable data. |
| Explanations: | | |
| \uparrow : statistically significant and relevant positive effect with low/unclear reliability of data | | |
| \downarrow : statistically significant and relevant negative effect with low/unclear reliability of data | | |
| $\uparrow\uparrow$: statistically significant and relevant positive effect with high reliability of data | | |
| $\downarrow \downarrow$: statistically significant and relevant negative effect with high reliability of data | | |
| \leftrightarrow : no statistically significant or relevant difference | | |
| arnothing: There are no usable data for the benefit assessment. | | |
| n.a.: not assessable | | |

b) Patients with 5q SMA type 2 and up to 3 copies of the SMN2 gene

No data available.

| Summary of results for relevant clinical endpoints | | |
|--|--|--|
| | | |

| Endpoint category | Direction of effect/ | Summary |
|--|----------------------|--------------------|
| | Risk of bias | |
| Mortality | Ø | No data available. |
| Morbidity | Ø | No data available. |
| Health-related quality | Ø | No data available. |
| of life | | |
| Side effects | Ø | No data available. |
| Explanations: | | |
| \uparrow : statistically significant and relevant positive effect with low/unclear reliability of data | | |
| \downarrow : statistically significant and relevant negative effect with low/unclear reliability of data | | |
| $\uparrow\uparrow$: statistically significant and relevant positive effect with high reliability of data | | |
| $\downarrow \downarrow$: statistically significant and relevant negative effect with high reliability of data | | |
| ↔: no statistically significant or relevant difference | | |
| arnothing: There are no usable data for the benefit assessment. | | |
| n.a.: not assessable | | |

c) Patients with 5q SMA type 3 and up to 3 copies of the SMN2 gene

No data available.

Summary of results for relevant clinical endpoints

| Endpoint category | Direction of effect/ Risk of bias | Summary |
|--|--------------------------------------|--------------------|
| Mortality | Ø | No data available. |
| Morbidity | Ø | No data available. |
| Health-related quality | Ø | No data available. |
| of life | | |
| Side effects | Ø | No data available. |
| Explanations: | | |
| \uparrow : statistically significant and relevant positive effect with low/unclear reliability of data | | |
| \downarrow : statistically significant and relevant negative effect with low/unclear reliability of data | | |
| $\uparrow\uparrow$: statistically significant and relevant positive effect with high reliability of data | | |
| $\downarrow \downarrow$: statistically significant and relevant negative effect with high reliability of data | | |
| \leftrightarrow : no statistically significant or relevant difference | | |
| arnothing: There are no usable data for the benefit assessment. | | |
| n.a.: not assessable | | |

d) Pre-symptomatic patients with 5q SMA and up to 3 copies of the SMN2 gene

No data available.

Summary of results for relevant clinical endpoints

| Endpoint category | Direction of effect/ Risk of bias | Summary |
|--|--------------------------------------|--------------------|
| Mortality | Ø | No data available. |
| Morbidity | Ø | No data available. |
| Health-related quality | Ø | No data available. |
| of life | | |
| Side effects | Ø | No data available. |
| Explanations: 个: statistically significant and relevant positive effect with low/unclear reliability of data | | |
| ↓: statistically significant and relevant negative effect with low/unclear reliability of data ↑↑: statistically significant and relevant positive effect with high reliability of data | | |
| $\downarrow \downarrow$: statistically significant and relevant negative effect with high reliability of data \leftrightarrow : no statistically significant or relevant difference | | |
| arnothing: There are no usable data for the benefit assessment. | | |
| n.a.: not assessable | | |

2. Number of patients or demarcation of patient groups eligible for treatment

- a) Patients with 5q SMA type 1
- b) Patients with 5q SMA type 2 and up to 3 copies of the SMN2 gene
- Patients with 5q SMA type 3 and up to 3 copies of the SMN2 gene approx. 45 to 65 patients
- Pre-symptomatic patients with 5q SMA and up to 3 copies of the SMN2 gene No specification possible

3. Requirements for a quality-assured application

The requirements in the product information are to be taken into account. The European Medicines Agency (EMA) provides the contents of the product information (summary of product characteristics, SmPC) for Zolgensma (active ingredient: onasemnogene abeparvovec) at the following publicly accessible link (last access: 11 August 2021):

https://www.ema.europa.eu/en/documents/product-information/zolgensma-epar-productinformation_en.pdf

The resolution of 20 November 2020 on requirements for a quality-assured application for the use of onasemnogene abeparvovec in spinal muscular atrophy provides further details.

This medicinal product was approved under "special conditions". This means that further evidence of the benefit of the medicinal product is anticipated. The European Medicines

Agency will evaluate new information on this medicinal product at a minimum once per year and update the product information where necessary.

4. Treatment costs

Treatment costs / Annual treatment costs:

Patient groups a, b and d

| Designation of the therapy | Treatment costs or annual treatment costs/patient |
|-------------------------------------|---|
| Medicinal product to be assessed: | |
| Onasemnogene abeparvovec | € 2,314,550 ¹ |
| Additionally required SHI services | different from patient to patient |
| Appropriate comparator therapy: | |
| Nusinersen 1st year | € 566,745.01 |
| additionally required SHI services: | non-quantifiable |
| Nusinersen subsequent years | € 261,574.62 |
| additionally required SHI services: | non-quantifiable |

Costs after deduction of statutory rebates (LAUER-TAXE®) as last revised: 15 October 2021)

Patient group c

| Designation of the therapy | Treatment costs or annual treatment costs/patient | |
|-------------------------------------|---|--|
| Medicinal product to be assessed: | | |
| Onasemnogene abeparvovec | € 2,314,550 ¹ | |
| Additionally required SHI services | different from patient to patient | |
| Best supportive care | different from patient to patient | |
| Appropriate comparator therapy: | | |
| Nusinersen 1st year | € 566,745.01 | |
| additionally required SHI services: | non-quantifiable | |
| Nusinersen subsequent years | € 261,574.62 | |
| additionally required SHI services: | non-quantifiable | |
| Best supportive care | different from patient to patient | |

Costs after deduction of statutory rebates (LAUER-TAXE®) as last revised: 15 October 2021)

¹ Single dose