

## **Kriterien zur Bestimmung der zweckmäßigen Vergleichstherapie**

**und**

**Recherche und Synopse der Evidenz zur Bestimmung der  
zweckmäßigen Vergleichstherapie nach § 35a SGB V**

**und**

**Schriftliche Beteiligung der wissenschaftlich-medizinischen  
Fachgesellschaften und der Arzneimittelkommission der  
deutschen Ärzteschaft (AkdÄ) zur Bestimmung der  
zweckmäßigen Vergleichstherapie nach § 35a SGB V**

**Vorgang: 2022-B-338 Luspatercept**

Stand: März 2023

## I. Zweckmäßige Vergleichstherapie: Kriterien gemäß 5. Kapitel § 6 VerfO G-BA

### Luspatercept

[Behandlung von Erwachsenen mit transfusionsabhängiger Anämie aufgrund von myelodysplastischen Syndromen (MDS)]

#### Kriterien gemäß 5. Kapitel § 6 VerfO

Sofern als Vergleichstherapie eine Arzneimittelanwendung in Betracht kommt, muss das Arzneimittel grundsätzlich eine Zulassung für das Anwendungsgebiet haben.	Siehe Übersicht „II. Zugelassene Arzneimittel im Anwendungsgebiet“.
Sofern als Vergleichstherapie eine nicht-medikamentöse Behandlung in Betracht kommt, muss diese im Rahmen der GKV erbringbar sein.	Die allogene Stammzelltransplantation kommt grundsätzlich als nicht-medikamentöse Behandlung im vorliegenden Anwendungsgebiet in Betracht. <sup>1</sup>
Beschlüsse/Bewertungen/Empfehlungen des Gemeinsamen Bundesausschusses zu im Anwendungsgebiet zugelassenen Arzneimitteln/nicht-medikamentösen Behandlungen	Beschlüsse über die Nutzenbewertung von Arzneimitteln mit neuen Wirkstoffen nach § 35a SGB V: <ul style="list-style-type: none"><li>• Luspatercept: Beschluss vom 21. Januar 2021</li></ul>
Die Vergleichstherapie soll nach dem allgemein anerkannten Stand der medizinischen Erkenntnisse zur zweckmäßigen Therapie im Anwendungsgebiet gehören.	Siehe systematische Literaturrecherche

<sup>1</sup> Zu dieser Behandlungsmethode wurde bisher keine Methodenbewertung durchgeführt.

## II. Zugelassene Arzneimittel im Anwendungsgebiet

Wirkstoff ATC-Code Handelsname	<b>Anwendungsgebiet</b> (Text aus Fachinformation)
Zu bewertendes Arzneimittel:	
Luspatercept B03XA06 Reblozyl	Zugelassenes Anwendungsgebiet: Reblozyl wird angewendet für die Behandlung von erwachsenen Patienten mit transfusionsabhängiger Anämie aufgrund von myelodysplastischen Syndromen (MDS) mit Ringsideroblasten, mit sehr niedrigem, niedrigem oder intermediärem Risiko, die auf eine Erythropoetin-basierte Therapie nicht zufriedenstellend angesprochen haben oder dafür nicht geeignet sind.
Azacitidin L01BC07 Vidaza	Vidaza ist angezeigt zur Behandlung von erwachsenen Patienten, die für eine Transplantation hämatopoetischer Stammzellen (HSZT) nicht geeignet sind und eines der folgenden Krankheitsbilder aufweisen: <ul style="list-style-type: none"> <li>- myelodysplastische Syndrome (MDS) mit intermediärem Risiko 2 oder hohem Risiko nach International Prognostic Scoring System (IPSS)</li> </ul>
Deferasirox V03AC03 Exjade	EXJADE ist auch angezeigt zur Behandlung der chronischen, transfusionsbedingten Eisenüberladung, wenn eine Deferoxamin-Therapie bei folgenden Patientengruppen kontraindiziert oder unangemessen ist: <ul style="list-style-type: none"> <li>- bei Kindern im Alter zwischen 2 und 5 Jahren mit Beta-Thalassämia major mit Eisenüberladung auf Grund häufiger Transfusionen (<math>\geq 7</math> ml/kg/Monat Erythrozytenkonzentrat),</li> <li>- bei Erwachsenen, Kindern und Jugendlichen im Alter von 2 Jahren oder älter mit Beta-Thalassämia major mit Eisenüberladung auf Grund seltener Transfusionen Transfusionen (<math>&lt; 7</math> ml/kg/Monat Erythrozytenkonzentrat),</li> <li>- <b>bei Erwachsenen, Kindern und Jugendlichen im Alter von 2 Jahren und älter mit anderen Anämien.</b></li> </ul>
Deferoxamin V03AC01 Desferal	Behandlung der chronischen Eisenüberladung, z. B. <ul style="list-style-type: none"> <li>- Transfusionshämosiderosen, insbesondere bei Thalassaemia major, sideroblastischer Anämie, autoimmunhämolytischer Anämie <b>und anderen chronischen Anämien</b>;</li> <li>- primärer (idiopathischer) Hämochromatose bei Patienten, deren Begleiterkrankungen (z. B. schwere Anämie, Herzerkrankungen, Hypoproteinämie) einen Aderlass ausschließen;</li> <li>- Eisenüberladung bei Patienten mit Porphyria cutanea tarda;</li> </ul> Behandlung der akuten Eisenvergiftung.

## II. Zugelassene Arzneimittel im Anwendungsgebiet

Epoetin alfa B03XA01 Erypo	ERYPO ist angezeigt zur Behandlung der symptomatischen Anämie (Hämoglobinspiegel $\leq 10$ g/dl) bei Erwachsenen mit primären Niedrigrisiko Myelodysplastischen Syndromen (MDS) (niedrig oder intermediär-1) und niedrigen Erythropoetin-Serumsiegeln (< 200 mU/ml).
Epoetin zeta B03XA0 Retacrit	Retacrit ist angezeigt zur Behandlung der symptomatischen Anämie (Hämoglobin-spiegel $\leq 10$ g/dl) bei Erwachsenen mit primären Niedrigrisiko Myelodysplastischen Syndromen (MDS) (niedrig oder intermediär-1) und niedrigen Erythropoetin-Serum-spiegeln (< 200 mU/ml).
Imatinib L01XE01 Glivec	Imatinib ist angezeigt zur Behandlung: - von Erwachsenen mit myelodysplastischen/myeloproliferativen Erkrankungen (MDS/MPD) in Verbindung mit Genumlagerungen des PDGF-Rezeptors (platelet-derived growth factor).
Lenalidomid L04AX04 Revlimid	Myelodysplastische Syndrome Revlimid als Monotherapie ist indiziert für die Behandlung von erwachsenen Patienten mit transfusionsabhängiger Anämie infolge myelodysplastischer Syndrome mit Niedrig- oder Intermediär-1-Risiko in Verbindung mit einer isolierten Deletion 5q als zytogenetische Anomalie, wenn andere Behandlungsoptionen nicht ausreichend oder nicht angemessen sind.
Erythrozyten- konzentrat n. a. n. a.	Anwendungsgebiete sind akute und chronische Anämien. Für die Indikation zur Erythrozytentransfusion lassen sich keine universell anwendbaren unteren Grenzwerte für Hämoglobin oder Hämatokrit festlegen. Die Ursache der Anämie soll möglichst geklärt werden und, falls möglich, eine kausale Therapie eingeleitet werden. Die Entscheidung für die Transfusion von Erythrozyten oder für eine andere, gleichwertige Therapie ist abhängig vom klinischen Gesamtzustand des Patienten.

Quellen: AMIice-Datenbank, Fachinformationen

## Abteilung Fachberatung Medizin

**Recherche und Synopse der Evidenz zur Bestimmung der zweckmäßigen Vergleichstherapie nach § 35a SGB V**

**Vorgang: 2022-B-338 (Luspatercept)**

Auftrag von: Abt. AM

Bearbeitet von: Abt. FB Med

Datum: 6. Februar 2023

## Inhaltsverzeichnis

Abkürzungsverzeichnis.....	3
1 Indikation .....	4
2 Systematische Recherche.....	4
3 Ergebnisse.....	5
3.1 Cochrane Reviews.....	5
3.2 Systematische Reviews .....	5
3.3 Leitlinien.....	5
4 Detaillierte Darstellung der Recherchestrategie.....	33
Referenzen .....	36

## Abkürzungsverzeichnis

AML	akute myeloische Leukämie
ATG	Antithymocyte globulin
AWMF	Arbeitsgemeinschaft der wissenschaftlichen medizinischen Fachgesellschaften
ECRI	ECRI Guidelines Trust
ESA	Erythropoiesis-stimulating agents
G-CSF	Granulocyte-colony stimulating factor
GIN	Guidelines International Network
GOR	Grade of Recommendations
GRADE	Grading of Recommendations, Assessment, Development and Evaluation
Hb	Haemoglobin
HCT	Hematopoietic Cell Transplantation
HCT-CI	Hematopoietic Cell Transplantation–Comorbidity Index
HgB	Hemoglobin
HMA	Hypomethylating agents
HSCT	hematopoietic stem cell transplantation
IPSS	International Prognostic Scoring System
IPSS-R	International Prognostic Scoring System
LDAC	Low-dose cytarabine
LoE	Level of Evidence
MAC	myeloablative conditioning
MDS	Myelodysplastische Syndrome
MDT	Multidisciplinary team meeting
NICE	National Institute for Health and Care Excellence
OS	Overall survival
RBC	Red blood cell
RIC	reduced-intensity conditioning
SIGN	Scottish Intercollegiate Guidelines Network
TPO	Thrombopoietin
TPO-RA	Thrombopoietin receptor agonists
TRIP	Turn Research into Practice Database
TRM	transplant-related mortality
WBC	White blood cell
WHO	World Health Organization

## 1 Indikation

Behandlung von erwachsenen Patientinnen und Patienten mit transfusionsabhängiger Anämie aufgrund von myelodysplastischen Syndromen (MDS)

*Hinweis zur Synopse: Informationen hinsichtlich nicht zugelassener Therapieoptionen sind über die vollumfängliche Darstellung der Leitlinienempfehlungen dargestellt.*

## 2 Systematische Recherche

Es wurde eine systematische Literaturrecherche nach systematischen Reviews, Meta-Analysen und evidenzbasierten systematischen Leitlinien zur Indikation myelodysplastisches Syndrom durchgeführt und nach PRISMA-S dokumentiert [A]. Die Recherchestrategie wurde vor der Ausführung anhand der PRESS-Checkliste begutachtet [B]. Es erfolgte eine Datenbankrecherche ohne Sprachrestriktion in: The Cochrane Library (Cochrane Database of Systematic Reviews), PubMed. Die Recherche nach grauer Literatur umfasste eine gezielte, iterative Handsuche auf den Internetseiten von Leitlinienorganisationen. Ergänzend wurde eine freie Internetsuche (<https://www.google.com/>) unter Verwendung des privaten Modus, nach aktuellen deutsch- und englischsprachigen Leitlinien durchgeführt.

Die Erstrecherche wurde am 31.05.2021 durchgeführt, die folgenden am 02.12.2021 und 11.01.2023. Die Recherchestrategie der Erstrecherche wurde unverändert übernommen und der Suchzeitraum jeweils auf die letzten fünf Jahre eingeschränkt. Die letzte Suchstrategie inkl. Angabe zu verwendeter Suchfilter ist am Ende der Synopse detailliert dargestellt. Die Recherchen ergaben insgesamt 465 Referenzen.

In einem zweistufigen Screening wurden die Ergebnisse der Literaturrecherche bewertet. Im ersten Screening wurden auf Basis von Titel und Abstract nach Population, Intervention, Komparator und Publikationstyp nicht relevante Publikationen ausgeschlossen. Zudem wurde eine Sprachrestriktion auf deutsche und englische Referenzen vorgenommen. Im zweiten Screening wurden die im ersten Screening eingeschlossenen Publikationen als Volltexte gesichtet und auf ihre Relevanz und methodische Qualität geprüft. Dafür wurden dieselben Kriterien wie im ersten Screening sowie Kriterien zur methodischen Qualität der Evidenzquellen verwendet. Basierend darauf, wurden insgesamt 4 Referenzen eingeschlossen. Es erfolgte eine synoptische Darstellung wesentlicher Inhalte der identifizierten Referenzen.

## 3 Ergebnisse

### 3.1 Cochrane Reviews

Es wurden keine relevanten Cochrane Reviews identifiziert.

### 3.2 Systematische Reviews

Es wurden keine relevanten systematischen Reviews identifiziert.

### 3.3 Leitlinien

---

**DeFilipp Z et al., 2022 [2].**

Hematopoietic Cell Transplantation in the Management of Myelodysplastic Syndrome: An Evidence-Based Review from the American Society for Transplantation and Cellular Therapy Committee on Practice Guidelines

#### Zielsetzung/Fragestellung

The goals of this update were to assemble and critically evaluate new evidence on the role of HCT in the treatment of adult patients with MDS, make treatment recommendations based on the available evidence, and identify areas of need for future research.

#### Methodik

##### Grundlage der Leitlinie

- Repräsentatives Gremium: unklar;
- Interessenkonflikte angegeben; finanzielle Unabhängigkeit nicht dargelegt;
- Systematische Suche; Auswahl und Bewertung der Evidenz unklar (siehe sonstige methodische Hinweise);
- Konsensusprozesse und externes Begutachtungsverfahren dargelegt;
- Empfehlungen der Leitlinie sind eindeutig und die Verbindung zu der zugrundeliegenden Evidenz ist explizit dargestellt;
- Regelmäßige Überprüfung der Aktualität gesichert: unklar.

##### Recherche/Suchzeitraum:

- Pubmed bis 01.07.2021

#### LoE

1++	High-quality meta-analyses, systematic reviews of randomized controlled trials (RCTs), or RCTs with a very low risk of bias
1+	Well-conducted meta-analyses, systematic reviews of RCTs, or RCTs with a low risk of bias Meta-analyses, systematic reviews of RCTs, or RCTs with a high risk of bias
1-	High-quality systematic reviews of case-control or cohort studies.
2++	High quality case-control or cohort studies with a very low risk of confounding, bias, or chance, and a high probability that the relationship is causal

2+	Well-conducted case-control or cohort studies with a low risk of confounding, bias, or chance, and a moderate probability that the relationship is causal
2 –	Case-control or cohort studies with a high risk of confounding, bias, or chance, and a significant risk that the relationship is not causal
3	Nonanalytic studies (eg, case reports, case series)
4	Expert opinion

## GoR

A	At least 1 meta-analysis, systematic review, or RCT rated as 1++ and directly applicable to the target population or a systematic review of RCTs or a body of evidence consisting principally of studies rated as 1+, directly applicable to the target population, and demonstrating overall consistency of results.
B	A body of evidence including studies rated as 2++, directly applicable to the target population, and demonstrating overall consistency of results or extrapolated evidence from studies rated as 1++ or 1+.
C	A body of evidence including studies rated as 2+, directly applicable to the target population, and demonstrating overall consistency of results or extrapolated evidence from studies rated as 2++.
D	Evidence level 3 or 4 or extrapolated evidence from studies rated as 2+.

## Sonstige methodische Hinweise

- In die Bestimmung des „Level of Evidence“ soll eine Einschätzung des Biasrisikos eingehen. Ergebnisse der Biasrisikobewertung sind der Publikation nicht zu entnehmen.

## Empfehlungen

Transplantation Indications and Patient-Specific Factors

Indication/Consideration	Recommendation	Grade of Recommendation	Highest Level of Evidence	References
Transplantation indications				
Should allogeneic HCT routinely be offered early for advanced (int-2/high) de novo MDS?	Yes	A	1++	7-12
Should allogeneic HCT routinely be offered early for lower risk (low/int-1) de novo MDS?	No	B	1+	7-9, 17-22
Patient-specific considerations				
Should eligibility for HCT in MDS be limited by age?	No	B	1++	10-12, 23-28
Should eligibility for HCT in MDS be limited by comorbidity?	Unclear	C	2+	23-28, 32

### Question 2: What is the role of allogeneic HCT in lower-risk MDS?

Patients with lower-risk MDS experience favorable outcomes after allogeneic HCT, largely owing to the lower risk of disease relapse. However, in the aforementioned comparison decision analyses, nontransplantation approaches consistently were associated with improved life expectancy for patients with lower-risk disease (low/int-1 by IPSS or very low/low by IPSS-R) [7-9,17]. Thus, most patients with lower-risk MDS who undergo HCT have a disease- or patient-related variable that the treating physician deems of high clinical concern. A limited number of studies have reported on transplantation outcomes for lower-risk MDS [18-22]. One prospective study randomized 182 subjects with IPSS-low, int-1, or int-2 MDS to HLA-matched HCT or nontransplantation therapies; 44 subjects (24%) had int-2 risk disease. The complete response rate was higher in the HCT group compared with the control group (53.8% versus 33%; P< .05); 4-year OS was also higher in the HCT group (79% versus 56%; P< .05). Of note, some nontransplantation therapies used in that study are not standard in North America [20]. In the largest retrospective study reported to date, a European Society for Blood and Marrow Transplantation (EBMT) registry analysis evaluated outcomes for 246 patients who underwent HCT for IPSS-low or int-1 risk disease [21]. Patients in that study were younger (<20% over age 61), and 76% were reclassified as intermediate or higher risk by IPSS-R. At 3 years, OS was 58% and the relapse rate was 16%.

Based on these data, we currently recommend that allogeneic HCT not be routinely offered to patients with lower-risk MDS (low or int-1 by IPSS) as an initial treatment strategy (Grade B, Level 1+). Most patients should be treated with nontransplantation approaches and undergo HCT only if disease progresses to have higher-risk features or fails to improve with nontransplantation therapies. However, there may be scenarios in which HCT is performed in IPSS lower-risk disease (eg, younger patients with transfusion-dependent disease despite best supportive care or those with multiple or adverse somatic mutations). Finally, investigation into the prognostic significance of somatic mutations, particularly in lower-risk MDS, is warranted and may modify future recommendations.

7. Cutler CS, Lee SJ, Greenberg P, et al. A decision analysis of allogeneic bone marrow transplantation for the myelodysplastic syndromes: delayed transplantation for low-risk myelodysplasia is associated with improved outcome. *Blood*. 2004;104:579–585.
8. Koreth J, Pidala J, Perez WS, et al. Role of reduced-intensity conditioning allogeneic hematopoietic stem-cell transplantation in older patients with de novo myelodysplastic syndromes: an international collaborative decision analysis. *J Clin Oncol*. 2013;31:2662–2670.
9. Alessandrino EP, Della Porta MG, Malcovati L, et al. Optimal timing of allogeneic hematopoietic stem cell transplantation in patients with myelodysplastic syndrome. *Am J Hematol*. 2013;88:581–588.
10. Robin M, Porcher R, Ades L, et al. HLA-matched allogeneic stem cell transplantation improves outcome of higher risk myelodysplastic syndrome a prospective study on behalf of SFGM-TC and GFM. *Leukemia*. 2015;29:1496–1501.
11. Kröger N, Sockel K, Wolschke C, et al. Comparison between 5-azacytidine treatment and allogeneic stem-cell transplantation in elderly patients with advanced MDS according to donor availability (VidazaAllo Study). *J Clin Oncol*. 2021;39:3318–3327.
12. Nakamura R, Saber W, Martens MJ, et al. Biologic assignment trial of reduced-intensity hematopoietic cell transplantation based on donor availability in patients 50–75 years of age with advanced myelodysplastic syndrome. *J Clin Oncol*. 2021;39:3328–3339.
13. Della Porta MG, Jackson CH, Alessandrino EP, et al. Decision analysis of allogeneic hematopoietic stem cell transplantation for patients with myelodysplastic syndrome stratified according to the revised International Prognostic Scoring System. *Leukemia*. 2017;31:2449–2457.
14. Lee SE, Kim YJ, Yahng SA, et al. Survival benefits from reduced-intensity conditioning in allogeneic stem cell transplantation for young lower-risk MDS patients without significant comorbidities. *Eur J Haematol*. 2011;87:510–520.
15. Díez Campelo M, Sanchez-Barba M, Gomez-García de Soria VG, et al. Results of allogeneic stem cell transplantation in the Spanish MDS registry: prognostic factors for low risk patients. *Leuk Res*. 2014;38:1199–1206.
16. Yu ZP, Ding JH, Sun AN, Ge Z, Chen BA, Wu DP. A randomized study comparing stem cell transplantation versus conventional therapy for low and intermediate-risk myelodysplastic syndromes patients. *Stem Cells Dev*. 2017;26:1132–1139.
17. Robin M, Porcher R, Zinke-Cerwenka W, et al. Allogeneic haematopoietic stem cell transplant in patients with lower risk myelodysplastic syndrome: a retrospective analysis on behalf of the Chronic Malignancy Working Party of the EBMT. *Bone Marrow Transplant*. 2017;52:209–215.
18. Novak P, Zabelina T, Wolschke C, Ayuk F, Christopeit M, Kröger N. Allogeneic stem cell transplantation for patients with lower-risk myelodysplastic syndrome. *Biol Blood Marrow Transplant*. 2020;26:2047–2052.
19. McClune BL, Weisdorf DJ, Pedersen TL, et al. Effect of age on outcome of reduced-intensity hematopoietic cell transplantation for older patients with acute myeloid leukemia in first complete remission or with myelodysplastic syndrome. *J Clin Oncol*. 2010;28:1878–1887.
20. Muffly L, Pasquini MC, Martens M, et al. Increasing use of allogeneic hematopoietic cell transplantation in patients aged 70 years and older in the United States. *Blood*. 2017;130:1156–1164.
21. Modi D, Deol A, Kim S, et al. Age does not adversely influence outcomes among patients older than 60 years who undergo allogeneic hematopoietic stem cell transplant for AML and myelodysplastic syndrome. *Bone Marrow Transplant*. 2017;52:1530–1536.
22. Itonaga H, Ishiyama K, Aoki K, et al. Increased opportunity for prolonged survival after allogeneic hematopoietic stem cell transplantation in patients aged 60–69 years with myelodysplastic syndrome. *Ann Hematol*. 2019;98:1367–1381.
23. Carre M, Porcher R, Finke J, et al. Role of age and Hematopoietic Cell Transplantation-Specific Comorbidity Index in myelodysplastic patients undergoing an allograft: a retrospective study from the Chronic Malignancies Working Party of the European Group for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant*. 2020;26:451–457.
24. Atallah E, Logan B, Chen M, et al. Comparison of patient age groups in transplantation for myelodysplastic syndrome: the Medicare Coverage with Evidence Development study. *JAMA Oncol*. 2020;6:486–493.
25. Penack O, Pczynski C, Mohty M, et al. Association of pre-existing comorbidities with outcome of allogeneic hematopoietic cell transplantation. A retrospective analysis from the EBMT. *Bone Marrow Transplant*. 2022;57:183–190.

Disease- and Transplantation-Related Factors

Factor	Recommendation	Grade of Recommendation	Highest Level of Evidence	References
Disease considerations				
Should HCT be offered for patients with therapy-related MDS?	Yes	B	2++	36-41
Should patients be assessed for chromosomal anomalies and somatic mutations prior to HCT?	Yes	B	2++	51-57, 60-65
Should HCT be offered for patients with high-risk cytogenetic or molecular disease?	Yes	B	2++	51-57, 60-65
Should patients with MDS receive disease-directed therapy prior to HCT?	Unclear	C	2++	66-75
Conditioning regimen				
Are RIC regimens an acceptable alternative for adults considered unfit for MAC regimens?	Yes	A	1++	73, 82-91
Should MAC be the preferred conditioning intensity in fit patients?	Unclear	A	1++	73, 82-91

36. Kröger N, Brand R, van Biezen A, et al. Risk factors for therapy-related myelodysplastic syndrome and acute myeloid leukemia treated with allogeneic stem cell transplantation. *Haematologica*. 2009;94:542–549.
37. Litzow MR, Tarima S, Perez WS, et al. Allogeneic transplantation for therapy-related myelodysplastic syndrome and acute myeloid leukemia. *Blood*. 2010;115:1850–1857.
38. Aldoss I, Pham A, Li SM, et al. Favorable impact of allogeneic stem cell transplantation in patients with therapy-related myelodysplasia regardless of TP53 mutational status. *Haematologica*. 2017;102:2030–2038.
39. Kida M, Usuki K, Uchida N, et al. Outcome and risk factors for therapy-related myeloid neoplasms treated with allogeneic stem cell transplantation in Japan. *Biol Blood Marrow Transplant*. 2020;26:1543–1551.
40. Metheny L, Callander NS, Hall AC, et al. Allogeneic transplantation to treat therapy-related myelodysplastic syndrome and acute myelogenous leukemia in adults. *Transplant Cell Ther*. 2021;27:923.e1–923.e12.
41. Itonaga H, Kida M, Hamamura A, et al. Outcome of therapy-related myelodysplastic syndrome and oligoblastic acute myeloid leukemia after allogeneic hematopoietic stem cell transplantation: a propensity score matched analysis. *Hematol Oncol*. 2022;40:752–762.
42. Onida F, Brand R, van Biezen A, et al. Impact of the International Prognostic Scoring System cytogenetic risk groups on the outcome of patients with primary myelodysplastic syndromes undergoing allogeneic stem cell transplantation from human leukocyte antigen-identical siblings: a retrospective analysis of the European Society for Blood and Marrow Transplantation-Chronic Malignancies Working Party. *Haematologica*. 2014;99:1582–1590.
43. Deeg HJ, Scott BL, Fang M, et al. Five-group cytogenetic risk classification, monosomal karyotype, and outcome after hematopoietic cell transplantation for MDS or acute leukemia evolving from MDS. *Blood*. 2012;120:1398–1408.
44. Scheid C, de Wreede L, van Biezen A, et al. Validation of the revised IPSS at transplant in patients with myelodysplastic syndrome/transformed acute myelogenous leukemia receiving allogeneic stem cell transplantation: a retrospective analysis of the EBMT chronic malignancies working party. *Bone Marrow Transplant*. 2017;52:1519–1525.
45. Ustun C, Trottier BJ, Sachs Z, et al. Monosomal karyotype at the time of diagnosis or transplantation predicts outcomes of allogeneic hematopoietic cell transplantation in myelodysplastic syndrome. *Biol Blood Marrow Transplant*. 2015;21:866–872.
46. van Gelder M, de Wreede LC, Schetelig J, et al. Monosomal karyotype predicts poor survival after allogeneic stem cell transplantation in chromosome 7 abnormal myelodysplastic syndrome and secondary acute myeloid leukemia. *Leukemia*. 2013;27:879–888.
47. Konuma T, Miyazaki Y, Uchida N, et al. Outcomes of allogeneic hematopoietic stem cell transplantation in adult patients with myelodysplastic syndrome harboring trisomy 8. *Biol Blood Marrow Transplant*. 2017;23:75–80.
48. Garderet L, Ziagkos D, van Biezen A, et al. Allogeneic stem cell transplantation for myelodysplastic syndrome patients with a 5q deletion. *Biol Blood Marrow Transplant*. 2018;24:507–513.
49. Bejar R, Stevenson KE, Caughey B, et al. Somatic mutations predict poor outcome in patients with myelodysplastic syndrome after hematopoietic stem-cell transplantation. *J Clin Oncol*. 2014;32:2691–2698.
50. Della Porta MG, Galli A, Bacigalupo A, et al. Clinical effects of driver somatic mutations on the outcomes of patients with myelodysplastic syndromes treated with allogeneic hematopoietic stem-cell transplantation. *J Clin Oncol*. 2016;34:3627–3637.

62. Lindsley RC, Saber W, Mar BG, et al. Prognostic mutations in myelodysplastic syndrome after stem-cell transplantation. *N Engl J Med.* 2017;376:536–547.
63. Yoshizato T, Nannya Y, Atsuta Y, et al. Genetic abnormalities in myelodysplasia and secondary acute myeloid leukemia: impact on outcome of stem cell transplantation. *Blood.* 2017;129:2347–2358.
64. Hamilton BK, Rybicki L, Hirsch C, et al. Mutation clonal burden and allogeneic hematopoietic cell transplantation outcomes in acute myeloid leukemia and myelodysplastic syndromes. *Bone Marrow Transplant.* 2019;54:1281–1286.
65. Nazha A, Hu ZH, Wang T, et al. A personalized prediction model for outcomes after allogeneic hematopoietic cell transplant in patients with myelodysplastic syndromes. *Biol Blood Marrow Transplant.* 2020;26:2139–2146.
66. Warlick ED, Cioc A, Defor T, Dolan M, Weisdorf D. Allogeneic stem cell transplantation for adults with myelodysplastic syndromes: importance of pretransplant disease burden. *Biol Blood Marrow Transplant.* 2009;15:30–38.
67. Alzahrani M, Power M, Abou Mourad Y, et al. Improving Revised International Prognostic Scoring System pre-allogeneic stem cell transplantation does not translate into better post-transplantation outcomes for patients with myelodysplastic syndromes: a single-center experience. *Biol Blood Marrow Transplant.* 2018;24:1209–1215.
68. Yahng SA, Kim M, Kim TM, et al. Better transplant outcome with pretransplant marrow response after hypomethylating treatment in higher risk MDS with excess blasts. *Oncotarget.* 2017;8:12342–12354.
69. Gerds AT, Gooley TA, Estey EH, Appelbaum FR, Deeg HJ, Scott BL. Pretransplantation therapy with azacitidine vs induction chemotherapy and posttransplantation outcome in patients with MDS. *Biol Blood Marrow Transplant.* 2012;18:1211–1218.
70. Damaj G, Duhamel A, Robin M, et al. Impact of azacitidine before allogeneic stem-cell transplantation for myelodysplastic syndromes: a study by the Societe Francaise de Greffe de Moelle et de Therapie-Cellulaire and the Groupe-Francophone des Myelodysplasies. *J Clin Oncol.* 2012;30:4533–4540.
71. Potter VT, Iacobelli S, van Biezen A, et al. Comparison of intensive chemotherapy and hypomethylating agents before allogeneic stem cell transplantation for advanced myelodysplastic syndromes: a study of the myelodysplastic syndrome subcommittee of the chronic malignancies working party of the European Society for Blood and Marrow Transplant Research. *Biol Blood Marrow Transplant.* 2016;22:1615–1620.
72. Kako S, Kanda Y, Kato J, et al. The bridge treatment selected at the decision for transplantation did not affect the outcomes in patients with MDS. *Hematol Oncol.* 2017;35:341–349.
73. Festuccia M, Deeg HJ, Gooley TA, et al. Minimal identifiable disease and the role of conditioning intensity in hematopoietic cell transplantation for myelodysplastic syndrome and acute myelogenous leukemia evolving from myelodysplastic syndrome. *Biol Blood Marrow Transplant.* 2016;22:1227–1233.
74. Hunter AM, Komrokji RS, Yun S, et al. Baseline and serial molecular profiling predicts outcomes with hypomethylating agents in myelodysplastic syndromes. *Blood Adv.* 2021;5:1017–1028.
75. Prebet T, Gore SD, Esterini B, et al. Outcome of high-risk myelodysplastic syndrome after azacitidine treatment failure. *J Clin Oncol.* 2011;29:3322–3327.
76. Martino R, Iacobelli S, Brand R, et al. Retrospective comparison of reduced-intensity conditioning and conventional high-dose conditioning for allogeneic hematopoietic stem cell transplantation using HLA-identical sibling donors in myelodysplastic syndromes. *Blood.* 2006;108:836–846.
77. Luger SM, Ringden O, Zhang MJ, et al. Similar outcomes using myeloablative vs reduced-intensity allogeneic transplant preparative regimens for AML or MDS. *Bone Marrow Transplant.* 2012;47:203–211.
78. Martino R, Henseler A, van Lint M, et al. Long-term follow-up of a retrospective comparison of reduced-intensity conditioning and conventional high-dose conditioning for allogeneic transplantation from matched related donors in myelodysplastic syndromes. *Bone Marrow Transplant.* 2017;52:1107–1112.
79. Kröger N, Iacobelli S, Franke GN, et al. Dose-reduced versus standard conditioning followed by allogeneic stem-cell transplantation for patients with myelodysplastic syndrome: a prospective randomized phase III study of the EBMT (RICMAC trial). *J Clin Oncol.* 2017;35:2157–2164.
80. Scott BL, Pasquini MC, Logan BR, et al. Myeloablative versus reduced intensity hematopoietic cell transplantation for acute myeloid leukemia and myelodysplastic syndromes. *J Clin Oncol.* 2017;35:1154–1161.
81. Jentzsch M, Döhrling C, Linke R, et al. Comparison of non-myeloablative and reduced-intensity allogeneic stem cell transplantation in older patients with myelodysplastic syndromes. *Am J Hematol.* 2019;94:1344–1352.
82. Bejanyan N, Zhang M, Bo-Subait K, et al. Myeloablative conditioning for allogeneic transplantation results in superior disease-free survival for acute myelogenous leukemia and myelodysplastic syndromes with low/ intermediate but not high disease risk index: a Center for International Blood and Marrow Transplant Research study. *Transplant Cell Ther.* 2021;27:68.e1–68.e9.
83. Scott BL, Pasquini MC, Fei M, et al. Myeloablative versus reduced-intensity conditioning for hematopoietic cell transplantation in acute myelogenous leukemia and myelodysplastic syndromes—long-term follow-up of the BMT CTN 0901 clinical trial. *Transplant Cell Ther.* 2021;27:483.e1–483.e6.

90. Beelen DW, Treischel R, Stelljes M, et al. Treosulfan or busulfan plus fludarabine as conditioning treatment before allogeneic haemopoietic stem cell transplantation for older patients with acute myeloid leukaemia or myelodysplastic syndrome (MC-FludT.14/L): a randomised, non-inferiority, phase 3 trial. *Lancet Haematol.* 2020;7:e28–e39.
91. Oran B, Ahn KW, Fretham C, et al. Fludarabine and melphalan compared with reduced doses of busulfan and fludarabine improve transplantation outcomes in older patients with myelodysplastic syndromes. *Transplant Cell Ther.* 2021;27: 921.e1-921.e10.

---

**Killick S et al., 2021 [3].**

British Society for Haematology guidelines for the management of adult myelodysplastic syndromes

**Zielsetzung/Fragestellung**

The objective of these guidelines is to provide healthcare professionals with clear guidance on the management of adult patients with MDS.

**Methodik**

Grundlage der Leitlinie

- Repräsentatives Gremium: Betroffene waren am Review der Leitlinie, allerdings nicht an deren Erstellung beteiligt;
- Interessenkonflikte und finanzielle Unabhängigkeit dargelegt;
- Systematische Suche; Auswahl und Bewertung der Evidenz unklar (siehe sonstige methodische Hinweise);
- Formale Konsensusprozesse nicht beschrieben, externes Begutachtungsverfahren dargelegt;
- Empfehlungen der Leitlinie sind eindeutig und die Verbindung zu der zugrundeliegenden Evidenz ist explizit dargestellt;
- Regelmäßige Überprüfung der Aktualität gesichert.

Recherche/Suchzeitraum:

- Pubmed from 2012 to December 2020

## LoE / GoR

The BSH Guidelines Committee uses the GRADE nomenclature for evaluating levels of evidence and assessing the strength of recommendations in all Guidance. Details are available at: <http://www.gradeworkinggroup.org/index.htm>

### **Quality of Evidence and criteria for assigning the quality of evidence**

The quality of evidence is graded as high (A), moderate (B), low (C) or very low (D).

Type of evidence	Randomized trial = high (A) Observational study = low (C) Any other evidence = very low (D)
Decrease* grade if	<ul style="list-style-type: none"> <li>• Serious or very serious limitation to study quality</li> <li>• Important inconsistency</li> <li>• Some or major uncertainty about directness</li> <li>• Imprecise or sparse data</li> <li>• High probability of reporting bias</li> </ul> <p>*Each quality criteria can reduce the quality by one or, if very serious, by two levels</p>
Increase grade if	<ul style="list-style-type: none"> <li>• Strong evidence of association—significant relative risk of &gt; 2 (&lt; 0.5) based on consistent evidence from two or more observational studies, with no plausible confounders (+1)</li> <li>• Very strong evidence of association—significant relative risk of &gt; 5 (&lt; 0.2) based on direct evidence with no major threats to validity (+2)</li> <li>• Evidence of a dose response gradient (+1)</li> <li>• All plausible confounders would have reduced the effect (+1)</li> </ul>

In general:

- (A) **High**: further research is very unlikely to change our confidence in the estimate of effect
- (B) **Moderate**: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate
- (C) **Low**: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate
- (D) **Very Low**: any estimate of effect is very uncertain

### **Strength of Recommendation**

**Strong (grade 1):** Strong recommendations are made if clinicians are certain that benefits do, or do not, outweigh risks and burdens. Grade 1 recommendations can be applied uniformly to most patients and words such as "recommend", "offer" and "should" are appropriate.

**Weak (grade 2):** Weak recommendations are made if clinicians believe that benefits and risks and burdens are finely balanced, or appreciable uncertainty exists about the magnitude of benefits and risks. In addition, clinicians are becoming increasingly aware of the importance of patient values and preferences in clinical decision making. When, across the range of patient values, fully informed patients are liable to make different choices, guideline panels should offer weak recommendations. Grade 2 recommendations require judicious application to individual patients and words such as "suggest" and "consider" are appropriate.

## Sonstige methodische Hinweise

- Laut Publikation basiert die Einschätzung der „quality of evidence“ auf einer Bewertung mittels GRADE. In diese Einschätzung fließt regelhaft eine Einschätzung des Biasrisikos ein. Ergebnisse der Biasrisikobewertung sind der Publikation nicht zu entnehmen.

## **Empfehlungen**

### Supportive care

Supportive care should be offered to all patients with MDS and symptomatic cytopenias (1A).

Red cell transfusions should be given to improve symptomatic anaemia (1A).

Policies for transfusion, including haemoglobin thresholds for red cell transfusion, should take clinical factors into consideration, including patient-related factors (1A).

Matching for Rh, K or additional antigens should be offered in line with current BSH Guidelines for patients expected to receive regular red cell transfusions (2C).

Local policies should be in place for the management of neutropenic sepsis (1A).

Patients with stable MDS not receiving intensive chemotherapy and without signs of bleeding should not be offered prophylactic platelet transfusions (1A).

TPO receptor agonists may be used to reduce bleeding events in thrombocytopenic patients with low or intermediate-1 risk MDS (1A).

#### *Management of anaemia with transfusion*

Red cell transfusion dependency is associated with decreased overall and leukaemia-free survival in MDS, and reduced quality of life (QoL).<sup>5–7</sup> Transfusion therapy is associated with well-recognised complications including risks of alloimmunisation.<sup>8,9</sup> Antibodies to Rh and K antigens appear the most common,<sup>10</sup> but the exact role and cost effectiveness of extended red cell phenotyping remains unknown and local practices vary.<sup>11</sup> Irradiated blood products are recommended after a stem cell transplant or treatment with antithymocyte globulin (ATG), in keeping with the current BSH Guidelines on the use of irradiated blood components.<sup>12</sup>

Although the severity of anaemia has a major impact on QoL in MDS patients,<sup>13</sup> the degree to which this may be ameliorated by different policies for red cell transfusion is not known. Clinicians may choose to apply a policy for red cell transfusion that is individualised and targeted to symptoms, although in practice specific haemoglobin (Hb) thresholds are often applied. A common haemoglobin threshold of around 80 g/l was identified by a UK national audit, a survey in Australia<sup>14</sup> and findings from the European MDS Registry (EUMDS).<sup>13</sup> The only randomised trial of transfusions in MDS patients compared two transfusion thresholds (80 g/l, to maintain Hb 85–100 g/l against 105 g/l, maintaining 110–125 g/l).<sup>15</sup> In an exploratory analysis, the five main QoL domains were improved for participants in the liberal compared to restrictive arm.

#### *Management of neutropenia and infection*

National Institute for Health and Care Excellence (NICE) has published guidelines for the prevention and management of neutropenic sepsis in cancer patients (CG151 published September 2012).<sup>16</sup> The use of prophylactic granulocyte-colony stimulating factor (G-CSF) may be considered in patients with recurrent infections who have low-risk MDS and may be used (with prophylactic antibiotics) to support the delivery of azacitidine in selected higher-risk patients.

Although a randomised centre study showed that in patients undergoing chemotherapy, posaconazole prevented invasive fungal infections more effectively than did either fluconazole or itraconazole and improved overall survival (OS),<sup>17</sup> there is no evidence to suggest that this should be routinely given to all patients with MDS. The American Society of Clinical Oncology and Infectious Diseases of America guidelines suggest that a mould-active triazole is recommended for patients who are at risk of profound, protracted neutropenia (defined as  $<0.1\ 9\ 10^9/l \geq 7\ days$ , or other risk factors).<sup>18</sup>

#### *Management of thrombocytopenia and bleeding*

There is common but variable practice of platelet transfusion in MDS. There are no similar studies in MDS, but a retrospective study in patients with stable chronic severe aplastic anaemia described a ‘no-prophylaxis’ platelet transfusion approach.<sup>19–21</sup> Avoiding unnecessary platelet transfusions in patients without signs of bleeding reduces the need for outpatient attendance improving QoL and may reduce the risk of platelet refractoriness. Patients with chronic thrombocytopenia presenting with bleeding of World Health Organisation (WHO) grade 2 or above should receive platelet transfusions.

Alternative agents to platelet transfusions include the antifibrinolytic drug tranexamic acid and should be considered as a symptomatic measure in mucous membrane bleeding in appropriate patients with MDS, although randomized trial evidence is lacking.<sup>22</sup>

Thrombopoietin receptor agonists (TPO-RA), specifically romiplostim and eltrombopag, have been evaluated in randomized placebo-controlled studies in both low-risk MDS and high-risk MDS (the latter in combination with either chemotherapy, hypomethylating agents or lenalidomide).<sup>23–29</sup> There were fewer bleeding episodes and fewer platelet transfusion episodes in the romiplostim arm in the Low/INT-1 study, although this study was halted prematurely because of concerns about increasing blast cell counts in patients receiving active drug.<sup>25</sup> A subsequent meta-analysis of several such studies did not find a significant difference in transformation to AML between intervention with TPO-RAs and placebo.<sup>30</sup> A


moderate reduction in bleeding events compared with placebo controls was noted, but with no improvement in mortality. Ongoing studies are evaluating the safety and efficacy of eltrombopag in Low/INT-1 MDS with severe thrombocytopenia (<30 9 109/l), and interim analysis has shown platelet responses in 47% of the eltrombopag group compared to 3% in the placebo group.<sup>31</sup> Although their use in high-risk MDS cannot be recommended, the results are promising for TPO-RA with platelet responses in low or intermediate-1 risk MDS (47–65%).<sup>24,31</sup> TPO-RA are not currently licenced for use in MDS and although these agents should ideally be accessed within clinical trials, the overall safety data now with longer follow-up is reassuring.

5. Sekeres MA, Maciejewski JP, List AF, Steensma DP, Artz A, Swern AS, et al. Perceptions of disease state, treatment outcomes, and prognosis among patients with myelodysplastic syndromes: results from an internet-based survey. *Oncologist*. 2011;16:904–11.
6. Malcovati L, Della Porta MG, Strupp C, Ambaglio I, Kuendgen A, Nachtkamp K, et al. Impact of the degree of anemia on the outcome of patients with myelodysplastic syndrome and its integration into the WHO classification-based Prognostic Scoring System (WPSS). *Haematologica*. 2011;96:1433–40.
7. Wood EM, Mc Quilten ZK. Outpatient transfusions for myelodysplastic syndromes. *Hematology Am Soc Hematol Educ Program*. 2020;2020:167–74.
8. Evers D, Middelburg RA, de Haas M, Zalpuri S, de Vooght KMK, van de Kerkhof D, et al. Red-blood-cell alloimmunisation in relation to antigens' exposure and their immunogenicity: a cohort study. *Lancet Haematol*. 2016;3:e284–e292.
9. Singhal D, Kutyna MM, Chhetri R, Wee LYA, Hague S, Nath L, et al. Red cell alloimmunization is associated with development of autoantibodies and increased red cell transfusion requirements in myelodysplastic syndrome. *Haematologica*. 2017;102:2021–9.
10. Lin Y, Sasin A, Wells RA, Lenis M, Mamedov A, Callum J, et al. Prophylactic RhCE and Kell antigen matching: impact on alloimmunization in transfusion-dependent patients with myelodysplastic syndromes. *Vox Sang*. 2017;112:79–86.
11. British Committee for Standards in Haematology, Milkins H, Berryman C, Cantwell J, Elliott C, Haggas C, et al. Guidelines for pre-transfusion compatibility procedures in blood transfusion laboratories. British Committee for Standards in Haematology. *Transfus Med*. 2013;23(1):3–35.
12. Foukaneli T, Kerr P, Bolton-Maggs PHB, Cardigan R, Coles A, Gennery A, et al. Guidelines on the use of irradiated blood components. *Br J Haematol*. 2020;191(5):701–24.
13. Stauder R, Yu GE, Koinig KA, Bagguley T, Fenaux P, Symeonidis A, et al. Health-related quality of life in lower-risk MDS patients compared with age- and sex-matched reference populations: a European Leukemia-Net study. *Leukemia*. 2018;32:1380–92.
14. Mo A, McQuilten ZK, Wood EM, Weinkove R. Red cell transfusion thresholds in myelodysplastic syndromes: a clinician survey to inform future clinical trials. *Intern Med J*. 2017;47:695–8.
15. Stanworth SJ, Killick S, McQuilten ZK, Karakantza M, Weinkove R, Smethurst H, et al. Red cell transfusion in outpatients with myelodysplastic syndromes: a feasibility and exploratory randomised trial. *Br J Haematol*. 2020;189(2):279–90.
16. Phillips R, Hancock B, Graham J, Bromham N, Jin H, Berendse S. Prevention and management of neutropenic sepsis in patients with cancer: summary of NICE guidance. *BMJ*. 2012;345:e5368.
17. Cornely OA, Maertens J, Winston DJ, Perfect J, Ullmann AJ, Walsh TJ, et al. Posaconazole vs. fluconazole or itraconazole prophylaxis in patients with neutropenia. *N Engl J Med*. 2007;356:348–59.
18. Taplitz RA, Kennedy EB, Bow EJ, Crews J, Gleason C, Hawley DK, et al. Antimicrobial prophylaxis for adult patients with cancer-related immunosuppression: ASCO and IDSA clinical practice guideline update. *J Clin Oncol*. 2018;36:3043–54.
19. Estcourt LJ, Birchall J, Allard S, Bassey SJ, Hersey P, Kerr JP, et al. Guidelines for the use of platelet transfusions. *Br J Haematol*. 2017;176:365–94.
20. Kaufman RM, Djulbegovic B, Gernsheimer T, Kleinman S, Tinmouth AT, Capocelli KE, et al. Platelet transfusion: a clinical practice guideline from the AABB. *Ann Intern Med*. 2015;162:205–13.
21. Sagmeister M, Oec L, Gmur J. A restrictive platelet transfusion policy allowing long-term support of outpatients with severe aplastic anemia. *Blood*. 1999;93:3124–6.
22. Desborough M, Estcourt LJ, Chaimani A, Doree C, Hopewell S, Trivella M, et al. Alternative agents versus prophylactic platelet transfusion for preventing bleeding in patients with thrombocytopenia due to chronic bone marrow failure: a network meta-analysis and systematic review. *Cochrane Database Syst Rev*. 2016;10(10):CD012055.
23. Dickinson M, Cherif H, Fenaux P, Mittelman M, Verma A, Portella MSO, et al. Azacitidine with or without eltrombopag for first-line treatment of intermediate- or high-risk MDS with thrombocytopenia. *Blood*. 2018;132:2629–38.
24. Fenaux P, Muus P, Kantarjian H, Lyons RM, Larson RA, Sekeres MA, et al. Romiplostim monotherapy in thrombocytopenic patients with myelodysplastic syndromes: long-term safety and efficacy. *Br J Haematol*. 2017;178:906–13.

25. Giagounidis A, Mufti GJ, Fenaux P, Sekeres MA, Szer J, Platzbecker U, et al. Results of a randomized, double-blind study of romiplostim versus placebo in patients with low/intermediate-1-risk myelodysplastic syndrome and thrombocytopenia. *Cancer*. 2014;120:1838–46.
26. Greenberg PL, Garcia-Manero G, Moore M, Damon L, Roboz G, Hu K, et al. A randomized controlled trial of romiplostim in patients with lower intermediate-risk myelodysplastic syndrome receiving decitabine. *Leuk Lymphoma*. 2013;54:321–8.
27. Kantarjian H, Fenaux P, Sekeres MA, Becker PS, Boruchov A, Bowen D, et al. Safety and efficacy of romiplostim in patients with lower-risk myelodysplastic syndrome and thrombocytopenia. *J Clin Oncol*. 2010;28:437–44.
28. Mittelman M, Platzbecker U, Afanasyev B, Grosicki S, Wong RSM, Anagnostopoulos A, et al. Eltrombopag for advanced myelodysplastic syndromes or acute myeloid leukaemia and severe thrombocytopenia (ASPIRE): a randomised, placebo-controlled, phase 2 trial. *Lancet Haematol*. 2018;5:e34–e43.
29. Wang ES, Lyons RM, Larson RA, Gandhi S, Liu D, Matei C, et al. A randomized, double-blind, placebo-controlled phase 2 study evaluating the efficacy and safety of romiplostim treatment of patients with low or intermediate-1 risk myelodysplastic syndrome receiving lenalidomide. *J Hematol Oncol*. 2012;5:71.
30. Dodillet H, Kreuzer KA, Monsef I, Skoetz N. Thrombopoietin mimetics for patients with myelodysplastic syndromes. *Cochrane Database Syst Rev*. 2017;9:CD009883.
31. Oliva EN, Alati C, Santini V, Poloni A, Molteni A, Niscola P, et al. Eltrombopag versus placebo for low-risk myelodysplastic syndromes with thrombocytopenia (EQoL-MDS): phase 1 results of a single-blind, randomised, controlled, phase 2 superiority trial. *Lancet Haematol*. 2017;4: e127–e136.

## Management of low-risk MDS

### Algorithm for the management of low-risk MDS



ATG, antithymocyte globulin; CSA, ciclosporin-A; ESA, erythropoiesis-stimulating agent; IPSS, International Prognostic Scoring System; MDS, myelodysplastic syndrome

Patients with IPSS Low and Intermediate-1 (or IPSS-R very low, low or intermediate with a score up to 3.5) MDS with symptomatic anaemia, or asymptomatic anaemia and Hb < 100 g/l and who fulfil the criteria for a high or intermediate predictive Nordic score for response should be considered for a trial of therapy with an ESA (1A).

For maximum benefit, ESA treatment should be started as soon as appropriate after diagnosis of MDS and before established transfusion dependence (for maximum benefit) (1B).

Patients should receive a maximum trial period of 24 weeks of therapy. This should comprise eight weeks at the starting dose of ESA, a further eight weeks at the higher doses, if required, and finally with the addition of G-CSF for a further eight weeks, before considering the patient to have failed ESA therapy (2B).

Patients achieving a complete or partial erythroid response by accepted criteria should continue on long-term therapy at the minimum dose of ESA required to maintain the response or until the response is lost (2B).

The haemoglobin concentration should not be allowed to rise above 120 g/l (2C).

#### *Erythropoiesis-stimulating agents*

It is only recently that randomised controlled trials for erythropoiesis-stimulating agents (ESAs) have been performed in the EU<sup>32,33</sup> and these have led to the European licence of EPO-a (Eprex), but not darbepoetin (Aranesp), for the treatment of symptomatic anaemia (haemoglobin ≤ 100 g/l) in adults with IPSS Low- or INT-1 primary MDS who have low serum EPO levels (<200 iu/l). There is a suggestion of survival advantage for responders to ESA therapy, especially if they are non-transfused prior to starting ESA,<sup>34–36</sup> and improvements in global QoL scores for responders.<sup>32,37,38</sup>

Who should be offered ESA therapy? ESA therapy is considered first-line standard of care for appropriately selected low-risk MDS patients who should have pretreatment variables that predict a response. The validated Nordic score, shown in Table I, has been widely used.<sup>37</sup> An alternative model is the ITACA scoring system.<sup>39</sup> As the Nordic model more effectively identifies likely non-responders, it remains the preferred model.

ESA therapy should be considered in patients with low or INT-1 IPSS (or IPSS-R very low, low or intermediate with a risk score of up to 3.5), in the context of symptomatic anaemia and Hb < 100 g/l. If patients are symptomatic from anaemia at a higher Hb, then starting an ESA is at the clinician's discretion. Patients should fulfil criteria predictive of response by the Nordic score (score 0–1). There are data to suggest that starting ESA therapy within six months of diagnosis improves response rates and delays the onset of transfusions (80 vs 35 months).<sup>34,40</sup> Patients with higher-risk MDS should not generally be considered for ESA therapy because of poor responses, short survival and the likely use of hypomethylating agents and stem cell transplantation, which require red cell transfusion support.

#### *Initial treatment*

Treatment should be initiated with EPO-a or darbepoetin alone in all patients. The recommended starting dose for EPO-a is 30 000–40 000 units subcutaneous once weekly for eight weeks ([mds-europe.eu](http://mds-europe.eu)).<sup>32,41</sup> If there is no response at eight weeks, the dose can be increased to a maximum dose of 60 000 units/week (divided over one or two doses) for a further eight weeks. Doses of >60 000 units/week are not supported by scientific evidence. The starting dose for darbepoetin should be 300 µg once every 14 days or 150 µg once every seven days ([mds-europe.eu](http://mds-europe.eu)).<sup>42,43</sup> This can be increased after eight weeks in non-responders to a maximum of 300 µg per week for a further eight weeks.<sup>44</sup> The starting dose in the randomised Phase 3 study<sup>33</sup> was 500 µg once every three weeks. However, 81% of patients had an increase in the dose to 500 µg every two weeks in the open-label period leading to a higher erythroid response. The starting dose of EPO-a or darbepoetin in low body weight with stable anaemia and always in the case of reduced renal function should be lower ([mds-europe.eu](http://mds-europe.eu)).

Finally, it is recommended that all patients receive incremental therapy with ESA alone for 16 weeks, as above, and G-CSF is then added to the higher dose in all non-responders for a final eight-week trial.<sup>45,46</sup> G-CSF should be given to approximately double the starting white cell count (WBC) if <1.5 × 10<sup>9</sup>/l, or keep the WBC in the range 6–10 × 10<sup>9</sup>/l. A starting dose of 300 µg per week or in 2/3 divided doses, rising to 300 µg three times per week in non-responders, is appropriate. However, the dosing regimen should be tailored to individual patients according to need and response. Response monitoring, criteria for response and long-term therapy. Response criteria for defining response<sup>37</sup> are as follows:

- Complete erythroid response: achievement of Hb > 115 g/l and transfusion independence.
- Partial erythroid response: >20 g/l increment in Hb and transfusion independence, but Hb remains <115 g/l.

Some patients may achieve potentially beneficial longer gaps between transfusions, although this is not a formally recognised response criterion.

The risk of thrombosis in MDS patients responding to darbepoetin has been estimated at 2%<sup>42</sup> and between 0.3 and 1.1% in meta-analysis.<sup>45</sup> However, in the randomised trial of EPO-a there were no grade 3–4 thrombo-embolic or stroke episodes in 85 treated patients.<sup>32</sup> In the darbepoetin randomized controlled trial,<sup>33,24</sup> weeks of darbepoetin produced no new safety signals and only one thromboembolic event (PE) in the darbepoetin group. Although the risk of thrombosis is low, it seems appropriate to temporarily interrupt ESA therapy if there is a rapid rise in haematocrit, or if the Hb rises above 120 g/l. Lower doses can then be introduced with careful monitoring of response parameters.

### *Response monitoring, criteria for response and long-term therapy*

Response criteria for defining response<sup>37</sup> are as follows:

- Complete erythroid response: achievement of Hb > 115 g/l and transfusion independence.
- Partial erythroid response: >20 g/l increment in Hb and transfusion independence, but Hb remains <115 g/l.

Some patients may achieve potentially beneficial longer gaps between transfusions, although this is not a formally recognised response criterion.

The risk of thrombosis in MDS patients responding to darbepoetin has been estimated at 2%<sup>42</sup> and between 0.3 and 1.1% in meta-analysis.<sup>45</sup> However, in the randomised trial of EPO-a there were no grade 3–4 thrombo-embolic or stroke episodes in 85 treated patients.<sup>32</sup> In the darbepoetin randomized controlled trial,<sup>33</sup> 24 weeks of darbepoetin produced no new safety signals and only one thromboembolic event (PE) in the darbepoetin group. Although the risk of thrombosis is low, it seems appropriate to temporarily interrupt ESA therapy if there is a rapid rise in haematocrit, or if the Hb rises above 120 g/l. Lower doses can then be introduced with careful monitoring of response parameters.

32. Fenaux P, Santini V, Spiriti MAA, Giagounidis A, Schlag R, Radinoff A, et al. A phase 3 randomized, placebo-controlled study assessing the efficacy and safety of epoetin-alpha in anemic patients with low-risk MDS. Leukemia. 2018;32:2648–58.
33. Platzbecker U, Symeonidis A, Oliva EN, Goede JS, Delforge M, Mayer J, et al. A phase 3 randomized placebo-controlled trial of darbepoetin alfa in patients with anemia and lower-risk myelodysplastic syndromes. Leukemia. 2017;31:1944–50.
34. Garelius HKG, Johnston WT, Smith AG, Park S, de Swart L, Fenaux P, et al. Erythropoiesis-stimulating agents significantly delay the onset of a regular transfusion need in nontransfused patients with lower-risk myelodysplastic syndrome. J Intern Med. 2017;281:284–99.
35. Jädersten M, Malcovati L, Dybedal I, Giovanni Della Porta M, Invernizzi R, Montgomery SM, et al. Erythropoietin and granulocyte-colony stimulating factor treatment associated with improved survival in myelodysplastic syndrome. J Clin Oncol. 2008;26:3607–13.
36. Park S, Grabar S, Kelaidi C, Beyne-Rauzy O, Picard F, Bardet V, et al. Predictive factors of response and survival in myelodysplastic syndrome treated with erythropoietin and G-CSF: the GFM experience. Blood. 2008;111:574–82.
37. Hellstrom-Lindberg E, Gulbrandsen N, Lindberg G, Ahlgren T, Dahl IM, Dybedal I, et al. A validated decision model for treating the anaemia of myelodysplastic syndromes with erythropoietin + granulocyte colony-stimulating factor: significant effects on quality of life. Br J Haematol. 2003;120:1037–46.
38. Nilsson-Ehle H, Birgegard G, Samuelsson J, Antunovic P, Astermark J, Garelius H, et al. Quality of life, physical function and MRI T2\* in elderly low-risk MDS patients treated to a haemoglobin level of >/=120 g/L with darbepoetin alfa +/- filgrastim or erythrocyte transfusions. Eur J Haematol. 2011;87:244–52.
39. Buckstein R, Balleari E, Wells R, Santini V, Sanna A, Salvetti C, et al. ITACA: a new validated international erythropoietic stimulating agent-response score that further refines the predictive power of previous scoring systems. Am J Hematol. 2017;92:1037–46.
40. Park S, Kelaidi C, Sapena R, Vassilieff D, Beyne-Rauzy O, Coiteux V, et al. Early introduction of ESA in low risk MDS patients may delay the need for RBC transfusion: a retrospective analysis on 112 patients. Leuk Res. 2010;34:1430–6.
41. Garypidou V, Verrou E, Vakalopoulou S, Perifanis V, Tziomalos K, Venizelos I. Efficacy of a single, weekly dose of recombinant erythropoietin in myelodysplastic syndromes. Br J Haematol. 2003;123:958.
42. Gabrilove J, Paquette R, Lyons RM, Mushtaq C, Sekeres MA, Tomita D, et al. Phase 2, single-arm trial to evaluate the effectiveness of darbepoetin alfa for correcting anaemia in patients with myelodysplastic syndromes. Br J Haematol. 2008;142:379–93.
43. Giraldo P, Nomdedeu B, Loscertales J, Requena C, de Paz R, Tormo M, et al. Darbepoetin alpha for the treatment of anemia in patients with myelodysplastic syndromes. Cancer. 2006;107:2807–16.
44. Mannone L, Gardin C, Quarre MC, Bernard JF, Vassilieff D, Ades L, et al. High-dose darbepoetin alpha in the treatment of anaemia of lower risk myelodysplastic syndrome results of a phase II study. Br J Haematol. 2006;133:513–9.
45. Park S, Fenaux P, Greenberg P, Mehta B, Callaghan F, Kim C, et al. Efficacy and safety of darbepoetin alpha in patients with myelodysplastic syndromes: a systematic review and meta-analysis. Br J Haematol. 2016;174:730–47.
46. Park S, Greenberg P, Yucel A, Farmer C, O'Neill F, De Oliveira BC, et al. Clinical effectiveness and safety of erythropoietin-stimulating agents for the treatment of low- and intermediate-1-risk myelodysplastic syndrome: a systematic literature review. Br J Haematol. 2019;184:134–60.

### Iron chelation in MDS

All suitable lower-risk patients (IPSS low and intermediate-1; IPSS-R low and very low) should be considered for iron chelation therapy at the time they have received 20 units of red cells, or when the ferritin is more than 1 000 Ig/l (1B).

Iron chelation therapy should be considered in patients prior to stem cell transplant, if time allows (2C).

Expert opinion is that deferasirox (although only licensed second line in MDS) is the drug of choice based on tolerability, compliance and mature safety data (2C).

Deferiprone is not routinely recommended in MDS (2C).

Iron chelation therapy with deferasirox should be stopped if the ferritin falls below 500 Ig/l and desferrioxamine should be stopped if the ferritin falls below 1 000 Ig/l (2C).

Patients with MDS are at risk of developing iron overload from transfusion of red cells where iron build-up is inevitable (one unit of red blood cells delivers 200–250 mg iron), and there is also increased intestinal absorption of iron driven by ineffective erythropoiesis,<sup>48</sup> mostly relevant to MDSRS. Excessive iron ultimately leads to secondary end organ damage and cardiac disease remains the main non-leukaemic cause of death in MDS.<sup>49,50</sup>

#### *Iron overload is associated with adverse outcome in MDS.*

Retrospective studies have shown that OS is significantly shorter in transfusion-dependent MDS patients either through cardiac deaths, hepatic cirrhosis<sup>51,50</sup> or increased leukaemic progression.<sup>50</sup> The European LeukemiaNet MDS Registry showed that the risk of death in transfusion-dependent patients with detectable labile plasma iron levels is independent of risk of disease progression.<sup>52</sup> Iron overload also increases transplant-related mortality in haematopoietic stem cell transplantation (HSCT) in MDS patients<sup>53</sup> and total transfusion burden implied a worse prognosis in a European Society for Blood and Marrow Transplantation (EBMT) study.<sup>54</sup>

#### *Measuring iron loading*

Routine estimations of iron loading can be made by serial monitoring of ferritin and tracking of red cell units transfused. However, there is little correlation between units transfused, or serum ferritin, and the degree of organ iron deposition.

Magnetic resonance imaging (MRI) for R2 (liver proton relaxation rate),<sup>55</sup> or cardiac and liver T2\* assessments<sup>56</sup> can be used to help quantify hepatic and cardiac iron loading and its impact on organ function.

#### *Iron chelation can improve natural history*

Effective iron chelation may improve haemopoiesis. The EPIC study<sup>57</sup> and the GIMEMA group<sup>58</sup> showed an International Working Group (IWG) erythroid response in 15–25% of patients although median response duration was only eight weeks in the EPIC study. Platelet and neutrophil responses were also reported. Desferrioxamine has been shown to lower cardiac iron assessed by MRI measurements<sup>59</sup> and deferasirox has been shown to improve alanine transaminase (ALT) levels.<sup>60</sup> A German registry study showed that chelation therapy improved survival in almost 200 transfused lower-risk MDS patients,<sup>61</sup> supported by prospective data from the EUMDS registry.<sup>62</sup> Furthermore, it is now accepted that iron chelation prior to HSCT in congenital anaemia can improve transplant-related mortality.<sup>53</sup> Although this is not yet proven to be the case in haematological neoplasms including MDS, a recent EBMT joint expert panel recommended chelation in patients who have received more than 20 units of blood prior to HSCT.<sup>63</sup>

#### *Choice of iron chelator*

Desferrioxamine remains the most efficient iron chelator available and is given subcutaneously in overnight infusions, which may decrease the labile iron pool. However, many patients find it uncomfortable and cumbersome, reporting QoL issues. Deferasirox and deferiprone are given orally and are generally well tolerated, although deferiprone is associated with agranulocytosis in around 4% of patients. Deferiprone should not be used routinely in patients with MDS, and only after careful consideration with a haematologist experienced in treating MDS. It should be undertaken with very careful monitoring (weekly blood counts), and should not be used where the baseline neutrophils are  $<1.5 \times 10^9/l$ . Deferasirox is the only iron chelator currently licensed for use in MDS patients with proven reduction in labile iron and improved haemopoiesis in some patients.<sup>57,64</sup>

#### *Discussion of recommendations*

Iron chelation in lower-risk MDS patients. It is recommended that all suitable lower-risk patients (IPSS low and intermediate-1; IPSS-R low and very low) should be considered for iron chelation therapy around the time they have received 20 units of red cells, or when the ferritin is more than 1 000 Ig/l. Patients should have ferritin levels measured every 12 weeks and have ophthalmological and auditory examinations before commencing therapy and annually while on treatment. Iron chelation with deferasirox should be stopped if the ferritin falls below 500 Ig/l and desferrioxamine should be stopped if the ferritin falls below 1 000 Ig/l.

#### *Iron chelation in higher-risk MDS patients*

Patients who are considered suitable for HSCT should have iron levels monitored and iron chelation therapy given prior to transplant, if time allows.

#### *Drug recommendations*

Deferasirox is only licensed second line (after desferrioxamine) for the treatment of chronic iron overload due to blood transfusions in patients with anaemia, such as MDS. However, real-world experience is that deferasirox is better tolerated, compliance is far superior and safety data are now mature. For these reasons, expert opinion is that deferasirox is the drug of choice for transfusion-related iron overload in patients with MDS. Desferrioxamine remains an option in those resistant to or intolerant of deferasirox. The two drugs may be combined in exceptional circumstances with heavy cardiac iron overload, but only under the supervision of a haematologist experienced in MDS treatment, although there are no data to support the combination.

There is no contra-indication to the use of iron chelation in combination with other disease-modulating treatments such as lenalidomide or azacitidine.

48. Weintraub LR, Conrad ME, Crosby WH. Regulation of the intestinal absorption of iron by the rate of erythropoiesis. *Br J Haematol.* 1965;11:432–8.
49. Della Porta MG, Malcovati L. Clinical relevance of extra-hematologic comorbidity in the management of patients with myelodysplastic syndrome. *Haematologica.* 2009;94:602–6.
50. Malcovati L, Porta MGD, Pascutto C, Invernizzi R, Boni M, Travaglino E, et al. Prognostic factors and life expectancy in myelodysplastic syndromes classified according to WHO criteria: a basis for clinical decision making. *J Clin Oncol.* 2005;23:7594–603.
51. Goldberg SL, Chen E, Corral M, Guo A, Mody-Patel N, Pecora AL, et al. Incidence and clinical complications of myelodysplastic syndromes among United States Medicare beneficiaries. *J Clin Oncol.* 2010;28:2847–52.
52. de Swart L, Reiniers C, Bagguley T, van Marrewijk C, Bowen D, Hellstrom-Lindberg E, et al. Labile plasma iron levels predict survival in patients with lower-risk myelodysplastic syndromes. *Haematologica.* 2018;103:69–79.
53. Großkatthöfer M, Güclü ED, Lawitschka A, Matthes-Martin S, Mann G, Minkov M, et al. Ferritin concentrations correlate to outcome of hematopoietic stem cell transplantation but do not serve as biomarker of graft-versus-host disease. *Ann Hematol.* 2013;2:1121–8.
54. Cremers EMP, van Biezen A, de Wreede LC, Scholten M, Vitek A, Finke J, et al. Prognostic pre-transplant factors in myelodysplastic syndromes primarily treated by high dose allogeneic hematopoietic stem cell transplantation: a retrospective study of the MDS subcommittee of the CMWP of the EBMT. *Ann Hematol.* 2016;95:1971–8.
55. St. Pierre TG, Clark PR, Chua-anusorn W, Fleming AJ, Jeffrey GP, Olynyk JK, et al. Noninvasive measurement and imaging of liver iron concentrations using proton magnetic resonance. *Blood.* 2005;105:855–61.
56. Anderson LJ, Holden S, Davis B, Prescott E, Charrier CC, Bunce NH, et al. Cardiovascular T2-star (T2\*) magnetic resonance for the early diagnosis of myocardial iron overload. *Eur Heart J.* 2001;22:2171–9.
57. Gattermann N, Finelli C, Della Porta M, Fenaux P, Stadler M, Guerci-Bresler A, et al. Hematologic responses to deferasirox therapy in transfusion-dependent patients with myelodysplastic syndromes. *Haematologica.* 2012;97:1364–71.
58. Angelucci E, Santini V, Di Tucci AA, Quaresmini G, Finelli C, Volpe A, et al. Deferasirox for transfusion-dependent patients with myelodysplastic syndromes: safety, efficacy, and beyond (GIMEMA MDS0306 Trial). *Eur J Haematol.* 2014;92:527–36.
59. Jensen PD, Jensen FT, Christensen T, Eiskjaer H, Baandrup U, Nielsen JL. Evaluation of myocardial iron by magnetic resonance imaging during iron chelation therapy with deferrioxamine: indication of close relation between myocardial iron content and chelatable iron pool. *Blood.* 2003;101:4632–9.
60. Gattermann N, Finelli C, Porta MD, Fenaux P, Ganser A, Guerci-Bresler A, et al. Deferasirox in iron-overloaded patients with transfusion-dependent myelodysplastic syndromes: results from the large 1-year EPIC study. *Leuk Res.* 2010;4:1143–50.

61. Neukirchen J, Fox F, Kundgen A, Nachtkamp K, Strupp C, Haas R, et al. Improved survival in MDS patients receiving iron chelation therapy – a matched pair analysis of 188 patients from the Dusseldorf MDS registry. *Leuk Res.* 2012;36:1067–70.
62. Hoeks M, Yu GE, Langemeijer S, Crouch S, de Swart L, Fenaux P, et al. Impact of treatment with iron chelation therapy in patients with lower-risk lower myelodysplastic syndromes participating in the European MDS registry. *Haematologica.* 2019;105(3):640–51.
63. de Witte T, Bowen D, Robin M, Malcovati L, Niederwieser D, Yakoub-Agha I, et al. Allogeneic hematopoietic stem cell transplantation for MDS and CMML: recommendations from an international expert panel. *Blood.* 2017;129:1753–62.
64. List AF, Baer MR, Steensma DP, Raza A, Esposito J, Martinez-Lopez N, et al. Deferasirox reduces serum ferritin and labile plasma iron in RBC transfusion-dependent patients with myelodysplastic syndrome. *J Clin Oncol.* 2012;30:2134–9.

### MDS associated with del(5q)

Patients with IPSS Low or INT-1 or IPSS-R with a score <3.5 and MDS with del(5q) and symptomatic anaemia and who fulfil the criteria for a high or intermediate predictive score for response, should be first considered for a trial of therapy with ESAs (1B).

For transfusion-dependent patients unsuitable for a trial of ESAs, and for non-responders and patients losing their response to ESAs, who have IPSS Low or INT-1 MDS with del(5q), consider treatment with lenalidomide 10 mg daily for 21 days repeated every 28 days after careful discussion with the patient about risk and benefit (1B).

Selected MDS patients with del(5q) and IPSS Low/INT-1 or IPSS-R with a score <3.5 may be candidates for allogeneic stem cell transplantation. These include lenalidomide-treated patients who fail to achieve transfusion independence, those losing their response, and patients with transfusion dependence not considered suitable for lenalidomide (2B).

Lenalidomide is not currently recommended for patients with del(5q) and bone marrow blasts >5% or multiple (complex) cytogenetic abnormalities in addition to del(5q) (neither of which fall into this diagnostic category) or patients with IPSS INT-2/high (2B).

MDS with isolated del(5q) is a distinct diagnostic entity that features macrocytic anaemia, normal or high platelet count, characteristic non-lobulated megakaryocytes and <5% bone marrow blasts. A single additional cytogenetic abnormality other than -7 or -7q is permitted within this diagnostic category. It is associated with female preponderance and has a relatively indolent natural history, with a median survival of six years in those with an IPSS score of 0.65. Independent predictors for OS include transfusion dependence, age and thrombocytopenia.<sup>66</sup>

Responses of patients with del(5q) MDS to ESA are inferior to that seen in low-risk MDS patients lacking del(5q) (39% vs 52%).<sup>67,68</sup> Nonetheless, given the established safety and efficacy data for ESA, ESA should be first-line therapy for symptomatic anaemia in lower-risk MDS patients with del(5q).

The MDS004 study compared lenalidomide with placebo in low and INT-1 transfusion-dependent MDS with del(5q); 58%, 42% and 6% of patients receiving lenalidomide 10 mg, 5 mg or placebo, respectively, achieved transfusion independence.<sup>69</sup> Cytogenetic responses were also seen in the lenalidomide treatment groups. Lenalidomide is licensed for transfusion-dependent low/INT-1 MDS with isolated del(5q) (with up to one abnormality other than -7/7q) and is recommended for NHS commissioning (NICE TA322) for such patients who have failed or are unresponsive to ESAs.

Concerns about the risk of progression to AML with lenalidomide have not been confirmed in retrospective studies,<sup>70,71</sup> post-MDS-004 study monitoring,<sup>72,73</sup> or a recent meta-analysis.<sup>74</sup> Rather, improved survival and reduced risk of transformation have been shown. Nonetheless, the MDS-004 study showed that progression to AML was 40% at five years compared to historically reported data of 20%. Follow-up studies have demonstrated that clonal evolution from existing or acquired TP53 mutations result in higher rates of AML transformation in del(5q) MDS patients.<sup>75–77</sup> However, some TP53-mutated cases with del(5q) have durable (2–3 years) responses to lenalidomide. Thus, TP53 mutation is not a contra-indication to lenalidomide therapy, but requires careful discussion and monitoring in this subgroup. Thromboprophylaxis should be considered on an individual basis.

Selected patients may be candidates for allogeneic stem cell transplantation. Indications include:

- Intolerance to or unsuitable for lenalidomide.
- Lenalidomide-treated patients who fail to achieve transfusion independence.
- Those with TP53 mutation.

- Those with clonal or overt progression.
- Those with bone marrow fibrosis.

66. Germing U, Lauseker M, Hildebrandt B, Symeonidis A, Cermak J, Fenaux P, et al. Survival, prognostic factors and rates of leukemic transformation in 381 untreated patients with MDS and del(5q): a multicenter study. *Leukemia*. 2012;26:1286–92.
67. Cheson BD, Greenberg PL, Bennett JM, Lowenberg B, Wijermans PW, Nimer SD, et al. Clinical application and proposal for modification of the International Working Group (IWG) response criteria in myelodysplasia. *Blood*. 2006;108:419–25.
68. Kelaidi C, Park S, Brechignac S, Mannone L, Vey N, Dombret H, et al. Treatment of myelodysplastic syndromes with 5q deletion before the lenalidomide era; the GFM experience with EPO and thalidomide. *Leuk Res*. 2008;32:1049–53.
69. Fenaux P, Giagounidis A, Selleslag D, Beyne-Rauzy O, Mufti G, Mittelman M, et al. A randomized phase 3 study of lenalidomide versus placebo in RBC transfusion-dependent patients with low-/intermediate-1-risk myelodysplastic syndromes with del5q. *Blood*. 2011;118:3765–76.
70. Ades L, Le Bras F, Sebert M, Kelaidi C, Lamy T, Dreyfus F, et al. Treatment with lenalidomide does not appear to increase the risk of progression in lower risk myelodysplastic syndromes with 5q deletion. A comparative analysis by the Groupe Francophone des Myelodysplasies. *Haematologica*. 2012;97:213–8.
71. Kuendgen A, Lauseker M, List AF, Fenaux P, Giagounidis AA, Brandenburg NA, et al. Lenalidomide does not increase AML progression risk in RBC transfusion-dependent patients with low- or intermediate-1-risk MDS with del(5q): a comparative analysis. *Leukemia*. 2013;27:1072–9.
72. List AF, Bennett JM, Sekeres MA, Skikne B, Fu T, Shammo JM, et al. Extended survival and reduced risk of AML progression in erythroid-responsive lenalidomide-treated patients with lower-risk del(5q) MDS.[Erratum appears in *Leukemia*. 2015 Dec;29(12):2452; PMID: 26648407]. *Leukemia*. 2015;2014(28):1033–40.
73. Sanchez-Garcia J, Del Canizo C, Lorenzo I, Nomdedeu B, Luno E, de Paz R, et al. Multivariate time-dependent comparison of the impact of lenalidomide in lower-risk myelodysplastic syndromes with chromosome 5q deletion. *Br J Haematol*. 2014;166:189–201.
74. Lian XY, Zhang ZH, Deng ZQ, He PF, Yao DM, Xu ZJ, et al. Efficacy and safety of lenalidomide for treatment of low-/intermediate-1-risk myelodysplastic syndromes with or without 5q deletion: a systematic review and meta-analysis. *PLoS One*. 2016;11:e0165948.
75. Jadersten M, Saft L, Smith A, Kulasekararaj A, Pomplun S, Gohring G, et al. TP53 mutations in low-risk myelodysplastic syndromes with del (5q) predict disease progression. *J Clin Oncol*. 2011;29:1971–9.
76. Lode L, Menard A, Flet L, Richebourg S, Loirat M, Eveillard M, et al. Emergence and evolution of TP53 mutations are key features of disease progression in myelodysplastic patients with lower-risk del(5q) treated with lenalidomide. *Haematologica*. 2018;103:e143–e146.
77. Scharenberg C, Giai V, Pellagatti A, Saft L, Dimitriou M, Jansson M, et al. Progression in patients with low- and intermediate-1-risk del(5q) myelodysplastic syndromes is predicted by a limited subset of mutations. *Haematologica*. 2017;102:498–508.

### Allogeneic haematopoietic stem cell transplant in MDS

All transplant-eligible MDS patients should be discussed with a transplant physician at a MDT both at diagnosis and at disease progression (2B).

Additional prognostic factors such as transfusion burden, depth of cytopenias, cytogenetics and BM fibrosis should be assessed when considering the optimal timing of transplant for lower-risk MDS patients (2B).

Higher-risk MDS patients with >10% blasts may be considered for cytoreductive therapy or hypomethylating agents prior to transplant (2B).

Up-front transplant may be considered in patients with 5–10% blasts with slowly progressive disease or in those with a hypocellular or fibrotic BM (2B).

Transplant is not routinely recommended for patients with TP53 mutation in association with a complex monosomal karyotype due to poor outcomes (2B).

Eligibility for transplant should be guided by HCT-CI and EBMT risk score (2B).

Performance status and age should be used to inform choice of myeloablative or reduced-intensity conditioning (2B).

All transplant-eligible MDS patients should be discussed with a transplant physician at a MDT, both at diagnosis and with disease progression. The decision to transplant should be made on a case-by-case basis, evaluating patient, donor and disease factors known to influence transplant outcomes.<sup>101</sup>

#### *Factors influencing timing and decision to transplant*

##### *Lower-risk MDS*

The optimal time to transplant patients with lower-risk MDS remains an area of debate. Early transplant for the lowest-risk patients is generally not recommended due to subsequent reduction in life expectancy.<sup>102–104</sup>

To help guide decision-making, particularly in the IPSS INT-1 group, the ELN/EBMT guidelines<sup>63</sup> recommend the use of other poor prognostic factors such as transfusion dependency ( $\geq 2$  units of blood per month), significant cytopenias, e.g. platelet count  $<30 \times 10^9/l$ , neutrophils  $<0.3 \times 10^9/l$ , or very poor prognostic cytogenetics. Transfusion dependence, elevated ferritin and labile plasma iron levels correlate with increased transplant-related mortality (TRM) in MDS patients following transplantation.<sup>50,105–107</sup>


Transplant should be considered once the patient becomes transfusion-dependent, before iron overload occurs. However, if there is a delay to transplant then iron chelation should strongly be considered.

Patients with progressive disease such as increasing blast cells or acquisition of adverse cytogenetic abnormalities should be considered for transplant.<sup>63</sup> Therapy failures, for example to ESAs or lenalidomide, convey a worse prognosis and should prompt consideration of transplantation.<sup>108,109</sup> Furthermore, patients with isolated del(5q) and an associated TP53 mutation have a worse prognosis and greater chance of failing lenalidomide therapy.<sup>75,77</sup> Such patients should be considered for transplantation early in their disease course.<sup>110</sup>

Patients with MDS and severe bone marrow (BM) fibrosis experience worse outcomes following HSCT compared with mild/moderate fibrosis, or those lacking fibrosis.<sup>111</sup> As such, the presence of BM fibrosis should prompt early transplant consideration, ideally prior to progression to severe fibrosis.

## Management of high-risk MDS

### Algorithm for the management of high-risk MDS



IPSS, international prognostic scoring system; IPSS-R, IPSS-revised; HSCT, haematopoietic stem cell transplant; PS, performance status. \*Where possible, patients should be offered entry into a clinical trial

### High-risk patients NOT eligible for allogeneic transplant

Patients requiring treatment should be considered for any appropriate clinical trial.

In fit older patients lacking an adverse karyotype, the options of therapy with a hypomethylating agent versus intensive chemotherapy should be carefully discussed.

Where intensive chemotherapy outside a clinical trial is planned, standard AML induction regimens should be used (2B).

Azacitidine is the preferred hypomethylating agent and is recommended as first-line therapy for patients ineligible for stem cell transplant with IPSS Intermediate-2 and high-risk MDS (IPSS-R Intermediate (score >3.5)/high/very high-risk groups) or AML with 20–30% blasts. Grade 1A (on the basis of a single randomised control trial).

The recommended dose of azacitidine is 75 mg/m<sup>2</sup> daily for seven consecutive days but a 5–2–2 schedule (with a two-days weekend gap) is acceptable where it is not practical to offer seven consecutive days and outcomes with the two schedules appear comparable (2B).

Outcomes of patients treated with azacitidine in routine clinical practice show a considerably shorter OS than the pivotal clinical trial (12.4–18.9 months compared to 24.5 months). Patients should be made aware of this.

Responding patients should continue azacitidine while their response is maintained (1A).

The decision to stop or continue azacitidine in patients who fail to achieve any response after six cycles, but who have stable disease, is dependent upon clinician and patient preference (2B).

Patients failing therapy with hypomethylating agents should be considered for any appropriate clinical trial.

#### *Intensive chemotherapy for patients ineligible for allogeneic HSCT*

For patients not eligible for transplantation, intensive AML-style chemotherapy can be used in an attempt to achieve disease response and improve survival. Patients should be entered into clinical trials where possible. The advantages of intensive chemotherapy are the QoL improvement if complete remission (CR) is achieved, and the small possibility of long-term disease-free survival.

There have been reported cases of long-term survival (>4 years) in patients with high-risk MDS and lacking an unfavourable karyotype.<sup>82</sup> However, older patients frequently have comorbidities, making intensive regimens less well tolerated. Overall, remission rates are lower (40–60%) than in de novo AML, remission duration is often shorter (median duration 10–12 months) and therapy-related complications of marrow aplasia (infection and haemorrhage) more frequent.<sup>82–85</sup>

Analysis of 160 patients over the age of 60 years with high-risk MDS or AML showed an early death rate of 10% and an inability to deliver consolidation chemotherapy in 40 of the 96 (42%) patients who achieved CR.<sup>84</sup> Compared to those with a normal karyotype who had a median survival of 18 months, those with a high-risk karyotype (involving 3 or more unrelated abnormalities or chromosome 7 abnormality) had a median survival of four months. The largest study of intensive chemotherapy for high-risk MDS broadly supports these data.<sup>86</sup> For this reason, it is recommended that cytogenetic results are available before committing to intensive chemotherapy in older patients with MDS, as there is no evidence to suggest this delay in treatment would be detrimental.<sup>87</sup>

#### *Disease-modifying agents in high-risk MDS*

Hypomethylating agents (azacitidine, decitabine) offer an alternative to intensive treatment in high-risk MDS. They are not curative but may result in transfusion independence, improved QoL and survival benefit and are well tolerated in the elderly and in patients with comorbidities.

Azacitidine is recommended by NICE and the Scottish Medicines Consortium as a treatment option for adult patients with MDS not eligible for HSCT (IPSS INT-2 or High) and for AML with 20–30% blasts and lineage dysplasia. The recommended dose is 75 mg/m<sup>2</sup> for seven consecutive days, repeated at 28-day intervals.

The AZA001 study<sup>88</sup> showed that azacitidine significantly increased OS compared to conventional-care regimens (median OS 24.5 vs 15.0 months).<sup>88</sup> Azacitidine also resulted in haematological responses; 45% of patients became transfusion-independent compared to 11% receiving conventional care. In a subgroup analysis of patients ≥75 years, azacytidine also significantly improved two-year OS compared to conventional care (55% vs 15%), suggesting that this is the treatment of choice in older higher-risk MDS patients with good performance status.<sup>89</sup>

Even patients with poor-prognosis cytogenetic profiles may benefit from azacitidine treatment.<sup>90</sup> Reliable molecular predictors of response have not been identified, although patients with poor-prognosis

indicators, including TP53 mutations, may respond. However, the presence of increasing numbers of mutations may be associated with a lower likelihood of response.<sup>91</sup>

Practical guidance for the delivery of azacitidine has been published.<sup>92</sup> Patients who receive less than six cycles or who fail to respond after six cycles have poor outcomes.<sup>93,94</sup> In the absence of progression and where azacitidine is tolerated, a minimum of six courses is recommended, with continued therapy for as long as response is maintained. Patients should have a marrow examination before starting treatment, after six courses (to assess response) and subsequently at clinician discretion should disease progression be suspected. In selected younger patients who achieve a CR with azacytidine and have good performance status, the option of HSCT should be re-visited.

Ongoing studies are exploring the combination of azacytidine with other agents in high-risk MDS.

The benefits of azacitidine have largely (but not uniformly) been confirmed in ‘real-world’ studies. However, OS in four large data sets has not matched that reported in the original pivotal trial.<sup>88</sup> The Canadian, Spanish and French Groups reported OS for azacitidine-treated patients with higher-risk MDS of 12.4, 13.4 and 13.5 months, respectively.<sup>93–95</sup>

Alternative dosing schedules for azacitidine include 75 mg/m<sup>2</sup> for five days, no treatment for two days, and two further days of treatment (5–2–2); 50 mg/m<sup>2</sup> on a 5–2–5 schedule or 75 mg/m<sup>2</sup> for five days.<sup>96</sup> In the Canadian real-world study of high-risk patients there was no difference in OS for patients treated with azacitidine for seven consecutive days compared with the 5–2–2 regimen,<sup>94</sup> and this is strongly preferred as the closest practical alternative if the licensed seven-day regimen is impractical.

Two Phase III studies comparing decitabine (15 mg/m<sup>2</sup> IV eight-hourly for three days every six weeks) with best supportive care in MDS have shown that some patients achieve CR, partial remission or haematological improvement. However, neither study showed significant improvement in OS.<sup>97,98</sup> In the ADOPT Phase II study of patients receiving decitabine 20 mg/m<sup>2</sup> for five days every four weeks,<sup>99</sup> CRs/marrow CRs of 32% and red cell (33%) and platelet (40%) transfusion independence were observed. Median survival was 19.4 months.

No prospective randomised studies comparing azacytidine with decitabine have been reported in intermediate-2/high-risk MDS. Azacitidine is the preferred agent, and the only one approved for use in the UK.

Although low-dose cytarabine (LDAC) has activity in high-risk MDS, the superiority of azacitidine over LDAC in the AZA001 study renders LDAC therapy obsolete in high-risk MDS.

Low-dose oral melphalan therapy could be considered for selective use in a rare group of patients, namely those with an excess of blasts (>5%) in a hypocellular marrow with a normal karyotype, for whom no alternative active therapy is available and/or appropriate. The majority of such patients will achieve CR with typical remission duration of 12 months.<sup>100</sup> Re-treatment will usually achieve a second remission but for a shorter duration. At melphalan-refractory relapse, patients are usually chemotherapy-resistant.

### Allogeneic haematopoietic stem cell transplant in MDS

All transplant-eligible MDS patients should be discussed with a transplant physician at a MDT both at diagnosis and at disease progression (2B).

Additional prognostic factors such as transfusion burden, depth of cytopenias, cytogenetics and BM fibrosis should be assessed when considering the optimal timing of transplant for lower-risk MDS patients (2B).

Higher-risk MDS patients with >10% blasts may be considered for cytoreductive therapy or hypomethylating agents prior to transplant (2B).

Up-front transplant may be considered in patients with 5–10% blasts with slowly progressive disease or in those with a hypocellular or fibrotic BM (2B).

Transplant is not routinely recommended for patients with TP53 mutation in association with a complex monosomal karyotype due to poor outcomes (2B).

Eligibility for transplant should be guided by HCT-CI and EBMT risk score (2B).

Performance status and age should be used to inform choice of myeloablative or reduced-intensity conditioning (2B).

All transplant-eligible MDS patients should be discussed with a transplant physician at a MDT, both at diagnosis and with disease progression. The decision to transplant should be made on a case-by-case basis, evaluating patient, donor and disease factors known to influence transplant outcomes.<sup>101</sup>

#### *Factors influencing timing and decision to transplant*

##### *Lower-risk MDS*

The optimal time to transplant patients with lower-risk MDS remains an area of debate. Early transplant for the lowest-risk patients is generally not recommended due to subsequent reduction in life expectancy.<sup>102–104</sup>

To help guide decision-making, particularly in the IPSS INT-1 group, the ELN/EBMT guidelines<sup>63</sup> recommend the use of other poor prognostic factors such as transfusion dependency ( $\geq 2$  units of blood per month), significant cytopenias, e.g. platelet count  $<30 \times 10^9/l$ , neutrophils  $<0.3 \times 10^9/l$ , or very poor prognostic cytogenetics. Transfusion dependence, elevated ferritin and labile plasma iron levels correlate with increased transplant-related mortality (TRM) in MDS patients following transplantation.<sup>50,105–107</sup>

Transplant should be considered once the patient becomes transfusion-dependent, before iron overload occurs. However, if there is a delay to transplant then iron chelation should strongly be considered.

Patients with progressive disease such as increasing blast cells or acquisition of adverse cytogenetic abnormalities should be considered for transplant.<sup>63</sup> Therapy failures, for example to ESAs or lenalidomide, convey a worse prognosis and should prompt consideration of transplantation.<sup>108,109</sup> Furthermore, patients with isolated del(5q) and an associated TP53 mutation have a worse prognosis and greater chance of failing lenalidomide therapy.<sup>75,77</sup> Such patients should be considered for transplantation early in their disease course.<sup>110</sup>

Patients with MDS and severe bone marrow (BM) fibrosis experience worse outcomes following HSCT compared with mild/moderate fibrosis, or those lacking fibrosis.<sup>111</sup> As such, the presence of BM fibrosis should prompt early transplant consideration, ideally prior to progression to severe fibrosis.

##### *Higher-risk MDS*

Early allogeneic HSCT offers a survival advantage in higher-risk MDS and suitable patients should be referred promptly to a transplant centre.<sup>102–104,112</sup> Inferior survival outcomes for patients with excess BM blasts ( $>5\%$ ) at the time of transplant have been reported.<sup>113</sup> It remains unclear, however, whether cytoreduction prior to transplant improves outcomes (regardless of BM blast percentage) over up-front transplantation.<sup>114</sup> In the absence of prospective data, patients with  $>10\%$  blasts may be considered for cytoreductive chemotherapy or hypomethylating agents (HMA) prior to transplant, particularly where immediate transplantation is not logistically possible.<sup>63</sup> Up-front transplantation should be considered where BM blasts are 5–10% in patients with slowly progressing disease, taking into account other patient- and disease-related factors. Patients with a hypocellular BM or presence of increased BM fibrosis with BM blasts up to 10% may also be considered for upfront transplant as prolonged cytopenia may occur with chemotherapy.

##### *Induction chemotherapy vs HMA prior to allogeneic HSCT*

Given the lack of available data from prospective, randomised trials, patients should be offered entry into a clinical trial, wherever possible. The ELN/EBMT support HSCT in suitable patients treated with HMA following attainment of CR.<sup>63</sup> However, emerging data from the VIDAZA ALLO study demonstrating early patient dropout due to treatment-related death or toxicity suggest that the number of HMA courses should be minimised.<sup>115</sup> For patients receiving induction chemotherapy, prolonged cytopenia may result; treatment should ideally be delivered once a donor has been identified (if a delay to commencing therapy is deemed acceptable).

Patients with a complex karyotype are more likely to exhibit TP53 mutation, contributing to their poor prognosis and therefore we recommend that all patients with complex karyotype are screened for TP53 mutation.<sup>110,116,117</sup> TP53 mutation is associated with resistance to conventional chemotherapy and early relapse.<sup>116,117</sup> In contrast, comparable response rates are observed following treatment with hypomethylating agents for MDS patients with TP53 mutation or wild-type TP53.<sup>118–121</sup> Patients with complex karyotype in the absence of TP53 mutation who require a reduction in blast count should be considered for clinical trials as there is no clear evidence to suggest whether intensive chemotherapy or HMA is better in this setting.<sup>122</sup>

##### *Choice of conditioning regimen*

The RICMAC trial showed no statistically significant difference in OS, relapse-free survival (RFS) or cumulative incidence of relapse at two years with reduced-intensity conditioning (RIC) or myeloablative conditioning (MAC).<sup>130</sup> A similar prospective trial demonstrated higher relapse rates for RIC versus MAC (48.3% vs 13.5%,  $P < 0.001$ ) leading to early trial closure.<sup>131</sup> In keeping with ELN/EBMT guidance, high-risk

patients with good performance status, lacking in comorbidity, may be candidates for MAC, reserving RIC for older, less fit patients.<sup>63</sup>

*Management of relapse post transplant*

Currently there are no standardised recommendations directing choice of therapy for relapse post HSCT and this is therefore not discussed further in this Guideline. Such patients may be best managed through accessing clinical trials where available.

50. Malcovati L, Porta MGD, Pascutto C, Invernizzi R, Boni M, Travaglino E, et al. Prognostic factors and life expectancy in myelodysplastic syndromes classified according to WHO criteria: a basis for clinical decision making. *J Clin Oncol.* 2005;23:7594–603.
63. de Witte T, Bowen D, Robin M, Malcovati L, Niederwieser D, Yakoub-Agha I, et al. Allogeneic hematopoietic stem cell transplantation for MDS and CMML: recommendations from an international expert panel. *Blood.* 2017;129:1753–62.
75. Jadersten M, Saft L, Smith A, Kulasekararaj A, Pomplun S, Gohring G, et al. TP53 mutations in low-risk myelodysplastic syndromes with del(5q) predict disease progression. *J Clin Oncol.* 2011;29:1971–9.
76. Lode L, Menard A, Flet L, Richebourg S, Loirat M, Eveillard M, et al. Emergence and evolution of TP53 mutations are key features of disease progression in myelodysplastic patients with lower-risk del(5q) treated with lenalidomide. *Haematologica.* 2018;103:e143–e146.
77. Scharenberg C, Gai V, Pellagatti A, Saft L, Dimitriou M, Jansson M, et al. Progression in patients with low- and intermediate-1-risk del(5q) myelodysplastic syndromes is predicted by a limited subset of mutations. *Haematologica.* 2017;102:498–508.
82. Wattel E, De Botton S, Luc Lai J, Preudhomme C, Lepelley P, Bauters F, et al. Long-term follow-up of de novo myelodysplastic syndromes treated with intensive chemotherapy: incidence of long-term survivors and outcome of partial responders. *Br J Haematol.* 1997;98:983–91.
83. Kantarjian H, O'Brien S, Cortes J, Giles F, Faderl S, Jabbour E, et al. Results of intensive chemotherapy in 998 patients age 65 years or older with acute myeloid leukemia or high-risk myelodysplastic syndrome: predictive prognostic models for outcome. *Cancer.* 2006c;106:1090–8.
84. Knipp S, Hildebrand B, Kundgen A, Giagounidis A, Kobbe G, Haas R, et al. Intensive chemotherapy is not recommended for patients aged >60 years who have myelodysplastic syndromes or acute myeloid leukemia with high-risk karyotypes. *Cancer.* 2007;110:345–52.
85. Morita Y, Kanamaru A, Miyazaki Y, Imanishi D, Yagasaki F, Tanimoto M, et al. Comparative analysis of remission induction therapy for highrisk MDS and AML progressed from MDS in the MDS200 study of Japan Adult Leukemia Study Group. *Int J Hematol.* 2010;91:97–103.
86. Kantarjian H, Beran M, Cortes J, O'Brien S, Giles F, Pierce S, et al. Long-term follow-up results of the combination of topotecan and cytarabine and other intensive chemotherapy regimens in myelodysplastic syndrome. *Cancer.* 2006;106:1099–109.
87. Sekeres MA, Elson P, Kalaycio ME, Advani AS, Copelan EA, Faderl S, et al. Time from diagnosis to treatment initiation predicts survival in younger, but not older, acute myeloid leukemia patients. *Blood.* 2009;113:28–36.
88. Fenaux P, Mufti GJ, Hellstrom-Lindberg E, Santini V, Finelli C, Giagounidis A, et al. Efficacy of azacitidine compared with that of conventional care regimens in the treatment of higher-risk myelodysplastic syndromes: a randomised, open-label, phase III study. *Lancet Oncol.* 2009;10:223–32.
89. Seymour JF, Fenaux P, Silverman LR, Mufti GJ, Hellstrom-Lindberg E, Santini V, et al. Effects of azacitidine compared with conventional care regimens in elderly (>/= 75 years) patients with higher-risk myelodysplastic syndromes. *Crit Rev Oncol Hematol.* 2010;76:218–27.
90. Ravandi F, Issa JP, Garcia-Manero G, O'Brien S, Pierce S, Shan J, et al. Superior outcome with hypomethylating therapy in patients with acute myeloid leukemia and high-risk myelodysplastic syndrome and chromosome 5 and 7 abnormalities. *Cancer.* 2009;115:5746–51.
91. Montalban-Bravo G, Takahashi K, Patel K, Wang F, Xingzhi S, Nogueras GM, et al. Impact of the number of mutations in survival and response outcomes to hypomethylating agents in patients with myelodysplastic syndromes or myelodysplastic/myeloproliferative neoplasms. *Oncotarget.* 2018;9:9714–27.
92. Fenaux P, Bowen D, Gattermann N, Hellstrom-Lindberg E, Hofmann WK, Pfeilstocker M, et al. Practical use of azacitidine in higher-risk myelodysplastic syndromes: an expert panel opinion. *Leuk Res.* 2010;34:1410–6.
93. Bernal T, Martinez-Camblor P, Sanchez-Garcia J, de Paz R, Luno E, Nomdedeu B, et al. Effectiveness of azacitidine in unselected high-risk myelodysplastic syndromes: results from the Spanish registry. *Leukemia.* 2015;29:1875–81.
94. Mozessohn L, Cheung MC, Fallahpour S, Gill T, Maloul A, Zhang L, et al. Azacitidine in the 'real-world': an evaluation of 1101 higher-risk myelodysplastic syndrome/low blast count acute myeloid leukaemia patients in Ontario, Canada. *Br J Haematol.* 2018;181:803–15.

95. Itzykson R, Thepot S, Quesnel B, Dreyfus F, Beyne-Rauzy O, Turlure P, et al. Prognostic factors for response and overall survival in 282 patients with higher-risk myelodysplastic syndromes treated with azacitidine. *Blood*. 2011;117:403–11.
96. Lyons RM, Cosgriff TM, Modi SS, Gersh RH, Hainsworth JD, Cohn AL, et al. Hematologic response to three alternative dosing schedules of azacytidine in patients with myelodysplastic syndromes. *J Clin Oncol*. 2009;27:1850–6.
97. Kantarjian H, Issa JP, Rosenfeld CS, Bennett JM, Albitar M, DiPersio J, et al. Decitabine improves patient outcomes in myelodysplastic syndromes: results of a phase III randomized study. *Cancer*. 2006b;106:1794–803.
98. Lubbert M, Suciu S, Baila L, Ruter BH, Platzbecker U, Giagounidis A, et al. Low-dose decitabine versus best supportive care in elderly patients with intermediate- or high-risk myelodysplastic syndrome (MDS) ineligible for intensive chemotherapy: final results of the randomized phase III study of the European Organisation for Research and Treatment of Cancer Leukemia Group and the German MDS Study Group. *J Clin Oncol*. 2011;29:1987–96.
99. Steensma DP, Baer MR, Slack JL, Buckstein R, Godley LA, Garcia-Manero G, et al. Multicenter study of decitabine administered daily for 5 days every 4 weeks to adults with myelodysplastic syndromes: the alternative dosing for outpatient treatment (ADOPT) trial. *J Clin Oncol*. 2009;27:3842–8.
100. Omoto E, Deguchi S, Takaba S, Kojima K, Yano T, Katayama Y, et al. Low-dose melphalan for treatment of high-risk myelodysplastic syndromes. *Leukemia*. 1996;10:609–14.
101. Shaffer BC, Ahn KW, Hu ZH, Nishihori T, Malone AK, Valcarcel D, et al. Scoring system prognostic of outcome in patients undergoing allogeneic hematopoietic cell transplantation for myelodysplastic syndrome. *J Clin Oncol*. 2016;34:1864–71.
102. Cutler CS, Lee SJ, Greenberg P, Deeg HJ, Perez WS, Anasetti C, et al. A decision analysis of allogeneic bone marrow transplantation for the myelodysplastic syndromes: delayed transplantation for low-risk myelodysplasia is associated with improved outcome. *Blood*. 2004;104:579–85.
103. Della Porta MG, Jackson CH, Alessandrino EP, Rossi M, Bacigalupo A, van Lint MT, et al. Decision analysis of allogeneic hematopoietic stem cell transplantation for patients with myelodysplastic syndrome stratified according to the revised International Prognostic Scoring System. *Leukemia*. 2017;31:2449–57.
104. Koreth J, Pidala J, Perez WS, Deeg HJ, Garcia-Manero G, Malcovati L, et al. Role of reduced-intensity conditioning allogeneic hematopoietic stem-cell transplantation in older patients with de novo myelodysplastic syndromes: an international collaborative decision analysis. *J Clin Oncol*. 2013;31:2662–70.
105. Alessandrino EP, Porta MG, Malcovati L, Jackson CH, Pascutto C, Bacigalupo A, et al. Optimal timing of allogeneic hematopoietic stem cell transplantation in patients with myelodysplastic syndrome. *Am J Hematol*. 2013;88:581–8.
106. Armand P, Kim HT, Cutler CS, Ho VT, Koreth J, Alyea EP, et al. Prognostic impact of elevated pretransplantation serum ferritin in patients undergoing myeloablative stem cell transplantation. *Blood*. 2007;109:4586–8.
107. Wermke M, Eckoldt J, Gotze KS, Klein SA, Bug G, de Wreede LC, et al. Enhanced labile plasma iron and outcome in acute myeloid leukaemia and myelodysplastic syndrome after allogeneic haemopoietic cell transplantation (ALLIVE): a prospective, multicentre, observational trial. *Lancet Haematol*. 2018;5:e201–e210.
108. Park S, Hamel JF, Toma A, Kelaidi C, Thepot S, Campelo MD, et al. Outcome of lower-risk patients with myelodysplastic syndromes without 5q deletion after failure of erythropoiesis-stimulating agents. *J Clin Oncol*. 2017;35:1591–7.
109. Jabbour EJ, Garcia-Manero G, Strati P, Mishra A, Al Ali NH, Padron E, et al. Outcome of patients with low-risk and intermediate-1-risk myelodysplastic syndrome after hypomethylating agent failure: a report on behalf of the MDS Clinical Research Consortium. *Cancer*. 2015;121:876–82.
110. Bejar R, Stevenson KE, Caughey B, Lindsley RC, Mar BG, Stojanov P, et al. Somatic mutations predict poor outcome in patients with myelodysplastic syndrome after hematopoietic stem-cell transplantation. *J Clin Oncol*. 2014;32:2691–8.
111. Kroger N, Zabelina T, van Biezen A, Brand R, Niederwieser D, Martino R, et al. Allogeneic stem cell transplantation for myelodysplastic syndromes with bone marrow fibrosis. *Haematologica*. 2011;96:291–7.
112. Kuendgen A, Strupp C, Aivado M, Hildebrandt B, Haas R, Gattermann N, et al. Myelodysplastic syndromes in patients younger than age 50. *J Clin Oncol*. 2006;24:5358–65.
113. Lim Z, Brand R, Martino R, van Biezen A, Finke J, Bacigalupo A, et al. Allogeneic hematopoietic stem-cell transplantation for patients 50 years or older with myelodysplastic syndromes or secondary acute myeloid leukemia. *J Clin Oncol*. 2010;28:405–11.
114. Schroeder T, Wegener N, Lauseker M, Rautenberg C, Nachtkamp K, Schuler E, et al. Comparison between upfront transplantation and different pretransplant cytoreductive treatment approaches in patients with high-risk myelodysplastic syndrome and secondary acute myelogenous leukemia. *Biol Blood Marrow Transplant*. 2019;25(8):1550–9.

115. Kroeger N, Sockel K, Wolschke C, Bethge W, Schlenk RF, Wolf D, et al. Prospective multicenter phase 3 study comparing 5-azacytidine (5-Aza) induction followed by allogeneic stem cell transplantation versus continuous 5-Aza according to donor availability in elderly MDS patients (55–70 years) (VidazaAllo study). *Blood.* 2018;132(Suppl. 1):208.
116. Rucker FG, Schlenk RF, Bullinger L, Kayser S, Teleanu V, Kett H, et al. TP53 alterations in acute myeloid leukemia with complex karyotype correlate with specific copy number alterations, monosomal karyotype, and dismal outcome. *Blood.* 2012;119:2114–21.
117. Sallman DA, Komrokji R, Vaupel C, Cluzeau T, Geyer SM, McGraw KL, et al. Impact of TP53 mutation variant allele frequency on phenotype and outcomes in myelodysplastic syndromes. *Leukemia.* 2016;30:666–73.
118. Bally C, Ades L, Renneville A, Sebert M, Eclache V, Preudhomme C, et al. Prognostic value of TP53 gene mutations in myelodysplastic syndromes and acute myeloid leukemia treated with azacitidine. *Leuk Res.* 2014;38:751–5.
119. Kulasekararaj AG, Smith AE, Mian SA, Mohamedali AM, Krishnamurthy P, Lea NC, et al. TP53 mutations in myelodysplastic syndrome are strongly correlated with aberrations of chromosome 5, and correlate with adverse prognosis. *Br J Haematol.* 2013;160:660–72.
120. Muller-Thomas C, Rudelius M, Rondak IC, Haferlach T, Schanz J, Huberle C, et al. Response to azacitidine is independent of p53 expression in higher-risk myelodysplastic syndromes and secondary acute myeloid leukemia. *Haematologica.* 2014;99:e179–e181.
121. Welch JS, Pettit AA, Miller CA, Fronick CC, O’Laughlin M, Fulton RS, et al. TP53 and decitabine in acute myeloid leukemia and myelodysplastic syndromes. *N Engl J Med.* 2016;375:2023–e36.
122. Damaj G, Duhamel A, Robin M, Beguin Y, Michallet M, Mohty M, et al. Impact of azacitidine before allogeneic stem-cell transplantation for myelodysplastic syndromes: a study by the Societe Francaise de Greffe de Moelle et de Therapie-Cellulaire and the Groupe-Francophone des Myelodysplasies. *J Clin Oncol.* 2012;30:4533–40.
130. Kroger N, Iacobelli S, Franke GN, Platzbecker U, Uddin R, Hubel K, et al. Dose-reduced versus standard conditioning followed by allogeneic stem-cell transplantation for patients with myelodysplastic syndrome: a prospective randomized phase III study of the EBMT (RICMAC Trial). *J Clin Oncol.* 2017;35:2157–64.
131. Scott BL, Pasquini MC, Logan BR, Wu J, Devine SM, Porter DL, et al. Myeloablative versus reduced-intensity hematopoietic cell transplantation for acute myeloid leukemia and myelodysplastic syndromes. *J Clin Oncol.* 2017;35:1154–61.

---

## Bohlius J et al., 2019 [1].

Management of cancer-associated anemia with erythropoiesis-stimulating agents:  
ASCO/ASH clinical practice guideline update

### Zielsetzung/Fragestellung

### Methodik

#### Grundlage der Leitlinie

- Repräsentatives Gremium;
- Interessenkonflikte und finanzielle Unabhängigkeit dargelegt;
- Systematische Suche, Auswahl und Bewertung der Evidenz;
- Formale Konsensusprozesse nicht beschrieben, externes Begutachtungsverfahren dargelegt;
- Empfehlungen der Leitlinie sind eindeutig und die Verbindung zu der zugrundeliegenden Evidenz ist explizit dargestellt;
- Regelmäßige Überprüfung der Aktualität gesichert: unklar.

#### Recherche/Suchzeitraum:

- PubMed and the Cochrane Library were searched for randomized controlled trials (RCTs) and meta-analyses of RCTs from January 31, 2010, through May 14, 2018

## LoE / GoR

- Recommendations reflect high, moderate, or low confidence that the recommendation reflects the net effect of a given course of action. The use of words like “must,” “must not,” “should,” and “should not” indicates that a course of action is recommended or not recommended for either most or many patients, but there is latitude for the treating physician to select other courses of action in individual cases.

## **Empfehlungen**

### Clinical question 1

To reduce the need for RBC transfusions, should ESAs be offered to patients who have chemotherapy-associated anemia?

#### Recommendation 1.1

Depending on clinical circumstances, ESAs may be offered to patients with chemotherapy-associated anemia whose cancer treatment is not curative in intent and whose HgB has declined to < 10 g/dL. RBC transfusion is also an option, depending on the severity of the anemia or clinical circumstances (Type: evidence based; Evidence quality: high; Strength of recommendation: strong).

#### Recommendation 1.2

ESAs should not be offered to patients with chemotherapy-associated anemia whose cancer treatment is curative in intent (Type: evidence based; Evidence quality: intermediate; Strength of recommendation: strong).

### Clinical question 2

To reduce the need for RBC transfusions, should ESAs be offered to anemic patients with cancer who are not receiving concurrent myelosuppressive chemotherapy?

#### Recommendation 2.1

ESAs should not be offered to most patients with nonchemotherapy-associated anemia (Type: informal consensus; Evidence quality: low; Strength of recommendation: strong).

#### Recommendation 2.2

ESAs may be offered to patients with lower-risk MDSs and a serum erythropoietin level ≤ 500 IU/L (Type: evidence based; Evidence quality: intermediate; Strength of recommendation: moderate).

In patients with MDS, one RCT evaluated the addition of epoetin beta to lenalidomide in 131 patients with RBC transfusion-dependent, low, or intermediate-1 risk (according to the International Prognostic Scoring System), ESA refractory, nondel(5q) MDS.<sup>26</sup> The combination of lenalidomide and epoetin beta increased the frequency of erythroid response relative to lenalidomide alone (39% v 23%; P = .04), but did not significantly affect duration of erythroid response (15 v 18 months; P = .64) or likelihood of transfusion independence (24% v 14%; P = .13). In subgroup analyses, patients with lower baseline serum erythropoietin levels had higher rates of erythroid response.

[...]

In patients with MDS, some studies suggest that patients with elevated baseline erythropoietin levels (> 500 IU/L) are unlikely to respond to ESA therapy.<sup>37</sup> Furthermore, a recent study has suggested that an even lower baseline erythropoietin level (< 200 IU/L) is associated with a better HgB response.<sup>38</sup> ESAs should be avoided in patients with MDS with elevated baseline erythropoietin levels (> 500 IU/L). Lower pretreatment RBC transfusion dependence (< 2 units per month) has also been associated with a higher likelihood of ESA response in patients with MDS.<sup>39</sup> Among the potential benefits of ESA therapy in patients

with MDS is avoidance of secondary hemochromatosis, particularly for lower risk patients who may have years of survival.

26. Toma A, Kosmider O, Chevret S, et al. Lenalidomide with or without erythropoietin in transfusion-dependent erythropoiesis-stimulating agent-refractory lower-risk MDS without 5q deletion. *Leukemia*. 2016;30:897-905.
37. Hellstrom-Lindberg E, Gulbrandsen N, Lindberg G, et al. A validated decision model for treating the anaemia of myelodysplastic syndromes with erythropoietin + granulocyte colony-stimulating factor: Significant effects on quality of life. *Br J Haematol*. 2003;120:1037-1046.
38. Fenaux P, Santini V, Spiriti MAA, et al. A phase 3 randomized, placebo-controlled study assessing the efficacy and safety of epoetin-a in anemic patients with low-risk MDS. *Leukemia*. 2018;32:2648-2658.
39. Hellstrom-Lindberg E, Ahlgren T, Beguin Y, et al. Treatment of anemia in myelodysplastic syndromes with granulocyte colony-stimulating factor plus erythropoietin: Results from a randomized phase II study and long-term follow-up of 71 patients. *Blood*. 1998;92:68-75.

### Clinical question 5

Among adult patients who receive an ESA for chemotherapy-associated anemia, do darbepoetin, epoetin beta and alfa originator, and currently available biosimilars of epoetin alfa differ with respect to safety or efficacy?

#### Recommendation 5

The Expert Panel considers epoetin beta and alfa, darbepoetin, and biosimilar epoetin alfa to be equivalent with respect to effectiveness and safety (Type: informal consensus; Evidence quality: intermediate; Strength of recommendation: moderate).

[...]

A retrospective study of patients with MDS and refractory anemia evaluated 46 patients treated with biosimilar epoetin alfa and 46 patients with originator epoetin alfa. Median time to reach an HgB level > 12 g/dL was 10.5 weeks (range, 3 to 16 weeks) among patients treated with the biosimilar and 12 weeks (range, 4 to 18 weeks) among patients treated with the originator product.<sup>31</sup>

[...]

31. Giordano G, Mondello P, Tambaro R, et al. Biosimilar epoetin a is as effective as originator epoetin-a plus liposomal iron (Sideral®), vitamin B12 and folates in patients with refractory anemia: A retrospective real-life approach. *Mol Clin Oncol*. 2015;3:781-784.

### Clinical question 6

Do ESAs increase the risk of thromboembolism?

#### Recommendation 6

ESAs increase the risk of thromboembolism, and clinicians should carefully weigh the risks of thromboembolism and use caution and clinical judgment when considering use of these agents (Type: evidence based; Evidence quality: high; Strength of recommendation: strong).

Meta-analyses and individual RCTs consistently report a 50%<sup>1,2</sup> to 75%<sup>16,25</sup> increased risk of thromboembolism and vascular arterial events among patients receiving ESA therapy. The Expert Panel continues to urge caution in the use of ESAs for patients judged to be at increased risk for venous thromboembolism. Several risk scores for predicting venous thromboembolism have been developed; these are discussed in more detail in the ASCO guideline on venous thromboembolism.<sup>42</sup> Special attention should be given to patients with multiple myeloma who are being treated with thalidomide or lenalidomide and doxorubicin or corticosteroids since they are at particularly increased thrombotic risk.<sup>43</sup> There are no data from RCTs investigating concomitant use of anticoagulants or aspirin to lessen this risk.

1. Grant MD, Piper M, Bohlius J, et al. AHRQ Comparative Effectiveness Reviews:Epoetin and Darbepoetin for Managing Anemia in Patients Undergoing Cancer Treatment: Comparative Effectiveness Update.

- Comparative Effectiveness Review No. 113 (AHRQ Publication No. 13-EHC077-EF). Rockville, MD: Agency for Healthcare Research and Quality; 2013. [www.effectivehealthcare.ahrq.gov/reports/final.cfm](http://www.effectivehealthcare.ahrq.gov/reports/final.cfm).
2. Tonia T, Mettler A, Robert N, et al. Erythropoietin or darbepoetin for patients with cancer. Cochrane Database Syst Rev. 2012;12:CD003407.
  16. Gao S, Ma JJ, Lu C. Venous thromboembolism risk and erythropoiesis-stimulating agents for the treatment of cancer-associated anemia: A meta-analysis. Tumour Biol. 2014;35:603-613.
  25. Zhan P, Wang Q, Qian Q, et al. Risk of venous thromboembolism with the erythropoiesis-stimulating agents (ESAs) for the treatment of cancer-associated anemia: A meta-analysis of randomized control trials. Linchuang Zhongliuxue Zazhi. 2012;1:19.
  42. Lyman GH, Khorana AA, Kuderer NM, et al. Venous thromboembolism prophylaxis and treatment in patients with cancer: American Society of Clinical Oncology clinical practice guideline update. J Clin Oncol. 2013;31:2189-2204.
  43. Bennett CL, Angelotta C, Yarnold PR, et al. Thalidomide- and lenalidomide-associated thromboembolism among patients with cancer. JAMA. 2006;296: 2558-2560.

### Clinical question 9

Among adult patients with chemotherapy-associated anemia who do not respond to ESA therapy (<1 to 2 g/dL increase in HgB or no decrease in transfusion requirements), does continuation of ESA therapy beyond 6 to 8 weeks provide a benefit?

#### Recommendation 9

ESAs should be discontinued in patients who do not respond within 6 to 8 weeks. Patients who do not respond to ESA treatment should be reevaluated for underlying tumor progression, iron deficiency, or other etiologies for anemia (Type: informal consensus; Evidence quality: intermediate; Strength of recommendation: strong).

Given the known harms of ESAs, the exposure to ESAs should be minimized. In patients who have received appropriate ESA dosing and who do not respond, ESAs should be stopped and not continued.

### Clinical question 10

Among adult patients with chemotherapy-associated anemia, does iron supplementation concurrent with an ESA reduce transfusion requirements?

#### Recommendation 10

Iron replacement may be used to improve HgB response and reduce RBC transfusions for patients receiving ESA with or without iron deficiency. Baseline and periodic monitoring of iron, total iron-binding capacity, transferrin saturation, or ferritin levels is recommended (Type: evidence based; Evidence quality: intermediate; Strength of recommendation: weak).

This recommendation changed from previous versions based on new information published after the last guideline. The Expert Panel believes that the use of iron supplementation in all patients receiving ESAs should be considered, independent of the iron status. This is based on evidence that iron supplementation reduces the risk for RBC transfusion. Additionally, in patients with evidence of iron deficiency, the cause of the deficiency should be investigated and corrected.

Oral and IV iron formulations are both acceptable options for iron supplementation. Choice of agents depends on patient and doctor preferences, formulation availability, cost, and comorbidities. IV iron preparations have the advantage of being able to deliver larger amounts of elemental iron in a single application and may also be more adequate in patients with poor oral intake or absorption problems. They have the disadvantages of being associated with more serious systemic reactions and higher costs. There is some limited evidence that IV iron is superior to oral iron based on improvement in HgB level. However, the results were not consistent across all other hematologic outcomes, and the quality of adverse outcomes reporting was poor.<sup>15,20,22</sup> Safety has been better studied in a systematic review and meta-analysis in patients with CKD, which showed no difference in mortality or serious adverse events in patients receiving intravenous iron, although there were more episodes of hypotension with IV iron.<sup>44</sup>

15. Gafter-Gvili A, Rozen-Zvi B, Vidal L, et al. Intravenous iron supplementation for the treatment of chemotherapy-induced anaemia—systematic review and meta-analysis of randomised controlled trials. *Acta Oncol.* 2013;52:18-29.
20. Mhaskar R, Wao H, Miladinovic B, et al. The role of iron in the management of chemotherapy-induced anemia in cancer patients receiving erythropoiesis-stimulating agents. *Cochrane Database Syst Rev.* 2016;2:CD009624.
22. Petrelli F, Borgonovo K, Cabiddu M, et al. Addition of iron to erythropoiesis-stimulating agents in cancer patients: A meta-analysis of randomized trials. *J Cancer Res Clin Oncol.* 2012;138:179-187.

**Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften), 2020 [4].**

S3-Leitlinie Supportive Therapie bei onkologischen PatientInnen; Langversion 1.3

### Leitlinienorganisation/Fragestellung

Therapieempfehlungen der Supportiven Therapie bei onkologischen Patienten

### Methodik

#### Grundlage der Leitlinie

- Repräsentatives Gremium;
- Interessenkonflikte und finanzielle Unabhängigkeit dargelegt;
- Systematische Suche, Auswahl und Bewertung der Evidenz;
- Formale Konsensusprozesse und externes Begutachtungsverfahren dargelegt;
- Empfehlungen der Leitlinie sind eindeutig und die Verbindung zu der zugrundeliegenden Evidenz ist explizit dargestellt;
- Regelmäßige Überprüfung der Aktualität gesichert.

#### LoE/GoR

- Schema der Evidenzgraduierung nach Oxford & GRADE

Tabelle 5: Schema der Empfehlungsgraduierung

Empfehlungsgrad	Beschreibung	Ausdrucksweise
A	Starke Empfehlung	Soll
B	Empfehlung	Sollte
O	Empfehlung offen	Kann

Tabelle 6: Konsensstärke

Konsensstärke	Prozentuale Zustimmung
Starker Konsens	> 95% der Stimmberchtigten
Konsens	> 75 – 95% der Stimmberchtigten
Mehrheitliche Zustimmung	> 50 – 75% der Stimmberchtigten
Dissens	< 50% der Stimmberchtigten

## Empfehlungen

### Myelodysplastisches Syndrom (MDS)

4.17.	Evidenzbasierte Empfehlung
<b>Empfehlungsgrad B</b>	Bei Patienten mit Myelodysplastischem Syndrom, die eine palliative Chemotherapie erhalten, sollte keine prophylaktische G-CSF Gabe erfolgen.
<b>Level of Evidence 1b</b>	Leitlinienadaptation: (Ossenkoppele, van der Holt et al. 1999, Zwierzina, Suciu et al. 2005, Vehreschild, Bohme et al. 2014)
<b>Abstimmung im Plenum</b>	Starker Konsens

### Hintergrund

In einer Studie mit 65 Hoch-Risiko MDS Patienten, die Daunorubicin ( $30 \text{ mg/m}^2$ , d1-3) und Cytarabin ( $200 \text{ mg/m}^2$ , d 1-7) erhielten, wurde der prophylaktische G-CSF Einsatz geprüft. Schwere Infektionen (WHO III-IV) traten häufiger im G-CSF Arm auf als im Kontrollarm (30% versus 19%). Die Inzidenz der FN wurde nicht berichtet (Ossenkoppele, van der Holt et al. 1999).

## 4 Detaillierte Darstellung der Recherchestrategie

Cochrane Library - Cochrane Database of Systematic Reviews (Issue 01 of 12, January 2023)  
am 09.01.2023

#	<b>Suchfrage</b>
1	[mh "Myelodysplastic Syndromes"]
2	(myelodysplas* OR dysmyelopoie* OR GATA2 NEXT deficienc* OR (GATA NEXT 2 NEXT deficienc*) OR PNH):ti,ab,kw
3	(an*emia* AND (refractory OR sideroblast*)):ti,ab,kw
4	((paroxysmal OR nocturnal OR h*emoly*) AND (h*emoglobinuria)):ti,ab,kw
5	(marchiafava* OR stru*bing* OR strübing*):ti,ab,kw
6	{OR #1-#5}
7	#6 with Cochrane Library publication date from Jan 2018 to present, in Cochrane Reviews

### Systematic Reviews in PubMed am 09.01.2023

verwendete Suchfilter:

*Konsentierter Standardfilter für Systematische Reviews (SR), Team Informationsmanagement der Abteilung Fachberatung Medizin, Gemeinsamer Bundesausschuss, letzte Aktualisierung am 02.01.2020.*

#	<b>Suchfrage</b>
1	myelodysplastic syndromes[mh]
2	myelodysplas*[tiab] OR dysmyelopoie*[tiab] OR GATA2 deficienc*[tiab] OR GATA-2 deficienc*[tiab] OR PNH[ti]
3	(anemia*[tiab] OR anaemia*[tiab]) AND (refractory[tiab] OR sideroblast*[tiab])
4	(paroxysmal[tiab] OR nocturnal[tiab] OR hemoly*[tiab] OR haemoly*[tiab]) AND (haemoglobinuria[tiab] OR hemoglobinuria[tiab])
5	marchiafava*[tiab] OR struebing*[tiab] OR strubing*[tiab] OR Strübing[tiab]
6	#1 OR #2 OR #3 OR #4 OR #5
7	(#6) AND (((Meta-Analysis[ptyp] OR systematic[sb] OR ((systematic review [ti] OR meta-analysis[pt] OR meta-analysis[ti] OR systematic literature review[ti] OR this systematic review[tw] OR pooling project[tw] OR (systematic review[tiab] AND review[pt]) OR meta synthesis[ti] OR meta-analy*[ti] OR integrative review[tw] OR integrative research review[tw] OR rapid review[tw] OR umbrella review[tw] OR consensus development conference[pt] OR practice guideline[pt] OR drug class reviews[ti] OR cochrane database syst rev[ta] OR acp journal club[ta] OR health technol assess[ta] OR evid rep technol assess summ[ta] OR jbi database system rev implement rep[ta]) OR (clinical guideline[tw] AND management[tw])) OR ((evidence based[ti] OR evidence-based medicine[mh] OR best practice*[ti] OR evidence synthesis[tiab]) AND (review[pt] OR diseases category[mh] OR behavior and behavior mechanisms[mh] OR therapeutics[mh] OR evaluation study[pt] OR validation study[pt] OR guideline[pt] OR pmcbook)) OR ((systematic[tw] OR systematically[tw] OR critical[tiab] OR (study selection[tw]) OR (predetermined[tw] OR inclusion[tw] AND criteri* [tw])) OR exclusion

#	Suchfrage
	criteri*[tw] OR main outcome measures[tw] OR standard of care[tw] OR standards of care[tw]) AND (survey[tiab] OR surveys[tiab] OR overview*[tw] OR review[tiab] OR reviews[tiab] OR search*[tw] OR handsearch[tw] OR analysis[ti] OR critique[tiab] OR appraisal[tw] OR (reduction[tw] AND (risk[mh] OR risk[tw])) AND (death OR recurrence))) AND (literature[tiab] OR articles[tiab] OR publications[tiab] OR publication [tiab] OR bibliography[tiab] OR bibliographies[tiab] OR published[tiab] OR pooled data[tw] OR unpublished[tw] OR citation[tw] OR citations[tw] OR database[tiab] OR internet[tiab] OR textbooks[tiab] OR references[tw] OR scales[tw] OR papers[tw] OR datasets[tw] OR trials[tiab] OR meta-analy*[tw] OR (clinical[tiab] AND studies[tiab])) OR treatment outcome[mh] OR treatment outcome[tw] OR pmcbook)) NOT (letter[pt] OR newspaper article[pt])) OR Technical Report[ptyp]) OR (((((trials[tiab] OR studies[tiab] OR database*[tiab] OR literature[tiab] OR publication*[tiab] OR Medline[tiab] OR Embase[tiab] OR Cochrane[tiab] OR Pubmed[tiab])) AND systematic*[tiab] AND (search*[tiab] OR research*[tiab])))) OR (((((((HTA[tiab]) OR technology assessment*[tiab])) OR technology report*[tiab]) OR (systematic*[tiab] AND review*[tiab]))) OR (systematic*[tiab] AND overview*[tiab])) OR meta-analy*[tiab]) OR (meta[tiab] AND analyz*[tiab])) OR (meta[tiab] AND analys*[tiab])) OR (meta[tiab] AND analyt*[tiab])))) OR (((review*[tiab]) OR overview*[tiab]) AND ((evidence[tiab]) AND based[tiab]))))))
8	(#7) AND ("2018/01/01"[PDAT] : "3000"[PDAT])
9	(#8) NOT "The Cochrane database of systematic reviews"[Journal]
10	(#9) NOT (retracted publication [pt] OR retraction of publication [pt])

### Leitlinien in PubMed am 09.01.2023

verwendete Suchfilter:

*Konsentierter Standardfilter für Leitlinien (LL), Team Informationsmanagement der Abteilung Fachberatung Medizin, Gemeinsamer Bundesausschuss, letzte Aktualisierung am 21.06.2017.*

#	Suchfrage
1	myelodysplastic syndromes[mh]
2	myelodysplas*[tiab] OR dysmyelopoie*[tiab] OR GATA2 deficienc*[tiab] OR GATA-2 deficienc*[tiab] OR PNH[ti]
3	(anemia*[tiab] OR anaemia*[tiab]) AND (refractory[tiab] OR sideroblast*[tiab])
4	(paroxysmal[tiab] OR nocturnal[tiab] OR hemoly*[tiab] OR haemoly*[tiab]) AND (haemoglobinuria[tiab] OR hemoglobinuria[tiab])
5	marchiafava*[tiab] OR struebing*[tiab] OR strubing*[tiab] OR Strübing[tiab]
6	#1 OR #2 OR #3 OR #4 OR #5
7	(#6) AND (Guideline[ptyp] OR Practice Guideline[ptyp] OR guideline*[Title] OR Consensus Development Conference[ptyp] OR Consensus Development Conference, NIH[ptyp] OR recommendation*[ti])
8	(#7) AND ("2018/01/01"[PDAT] : "3000"[PDAT])
9	(#8) NOT (retracted publication [pt] OR retraction of publication [pt])

### **Iterative Handsuche nach grauer Literatur, abgeschlossen am 11.01.2023**

- Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF)
- Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft, Deutsche Krebshilfe, AWMF)
- Nationale VersorgungsLeitlinien (NVL)
- National Institute for Health and Care Excellence (NICE)
- Scottish Intercollegiate Guideline Network (SIGN)
- World Health Organization (WHO)
- Alberta Health Service (AHS)
- European Society for Medical Oncology (ESMO)
- National Comprehensive Cancer Network (NCCN)
- National Cancer Institute (NCI)
- Dynamed / EBSCO
- Guidelines International Network (GIN)
- Trip Medical Database

## Referenzen

1. **Bohlius J, Bohlke K, Castelli R, Djulbegovic B, Lustberg MB, Martino M, et al.** Management of cancer-associated anemia with erythropoiesis-stimulating agents: ASCO/ASH clinical practice guideline update. *Blood Adv* 2019;3(8):1197-1210.
2. **DeFilipp Z, Ciurea SO, Cutler C, Robin M, Warlick ED, Nakamura R, et al.** Hematopoietic cell transplantation in the management of myelodysplastic syndrome: an evidence-based review from the American Society for Transplantation and Cellular Therapy Committee on practice guidelines. *Transplant Cell Ther* 2022 [Online ahead of print].
3. **Killick SB, Ingram W, Culligan D, Enright H, Kell J, Payne EM, et al.** British Society for Haematology guidelines for the management of adult myelodysplastic syndromes. *Br J Haematol* 2021;194(2):267-281.
4. **Leitlinienprogramm Onkologie (Deutsche Krebsgesellschaft (DKG), Deutsche Krebshilfe (DKH), Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften (AWMF))**. Supportive Therapie bei onkologischen PatientInnen; S3-Leitlinie, Langversion 1.3 [online]. AWMF-Registernummer 032-054OL. Berlin (GER): Leitlinienprogramm Onkologie; 2020. [Zugriff: 11.01.2023]. URL: [https://www.leitlinienprogramm-onkologie.de/fileadmin/user\\_upload/Downloads/Leitlinien/Supportivtherapie/LL\\_Supportiv\\_Langversion\\_1.3.pdf](https://www.leitlinienprogramm-onkologie.de/fileadmin/user_upload/Downloads/Leitlinien/Supportivtherapie/LL_Supportiv_Langversion_1.3.pdf).

- 
- [A] **Rethlefsen ML, Kirtley S, Waffenschmidt S, Ayala AP, Moher D, Page MJ, et al.** PRISMA-S: an extension to the PRISMA Statement for Reporting Literature Searches in Systematic Reviews. *Syst Rev* 2021;10(1):39. <https://doi.org/10.1186/s13643-020-01542-z>
- [B] **McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C**. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol* 2016;75:40-46. <https://doi.org/10.1016/j.jclinepi.2016.01.0>

**Beteiligung von Fachgesellschaften und der AkdÄ zu Fragen der Vergleichstherapie nach §35a Abs. 7 SGB V i.V.m. VerFO 5. Kapitel § 7 Abs. 6**

Verfahrens-Nr.: 2022-B-338

<b>Verfasser</b>	
DGHO	
Datum der Erstellung	22. Februar 2023

<b>Indikation</b>	
Behandlung von erwachsenen Patienten mit transfusionsabhängiger Anämie aufgrund von myelodysplastischen Syndromen (MDS) mit Ringsideroblasten, mit sehr niedrigem, niedrigem oder intermediärem Risiko, die auf eine Erythropoetin-basierte Therapie nicht zufriedenstellend angesprochen haben oder dafür nicht geeignet sind	
<b>Fragen zur Vergleichstherapie</b>	
Was ist der Behandlungsstandard in o.g. Indikation unter Berücksichtigung der vorliegenden Evidenz? Wie sieht die Versorgungspraxis in Deutschland aus?	
<b>Zusammenfassung</b>	
<p>MyeloDysplaStische Neoplasien (MDS, seit der WHO-Klassifikation 2022 wird der Begriff „Myelodysplastische Neoplasien“ verwendet [1], allerdings die Abkürzung „MDS“ beibehalten) sind klonale Erkrankungen der hämatopoetischen Stammzelle, die durch Dysplasien von Blut- und Knochenmarkzellen mit hämatopoetischer Insuffizienz und erhöhtem Risiko der Entwicklung einer akuten myeloischen Leukämie (AML) gekennzeichnet sind [2].</p> <p>Die Indikation zum Einsatz von Erythropoese-stimulierenden Faktoren bei Pat. mit MDS orientiert sich am Nordic Score. Kontraindikationen zum Einsatz von Erythropoese-stimulierenden Faktoren sind eine schwer einstellbare, arterielle Hypertonie und eine Überempfindlichkeit gegen den Wirkstoff oder sonstige Bestandteile.</p> <p>Standard in der Behandlung von erwachsenen Patienten mit transfusionsabhängiger Anämie aufgrund von myelodysplastischen Neoplasien (MDS) mit Ringsideroblasten, mit sehr niedrigem, niedrigem oder intermediärem Risiko, die auf eine Erythropoetin-basierte Therapie nicht zufriedenstellend angesprochen haben oder dafür nicht geeignet sind, ist die Gabe von Luspatercept.</p>	
<b>Fragestellung</b>	
Eine aktuelle Beratung für Studien und/oder für die Nutzenbewertung neuer Arzneimittel muss die geänderte WHO Klassifikation von 2022 berücksichtigen.	

## Stand des Wissens

Die MDS zählen mit einer Inzidenz von ca. 4-5/100.000 Einwohnern pro Jahr zu den häufigsten malignen hämatologischen Erkrankungen [3]. Im Alter über 70 Jahre steigt die Inzidenz auf >30/100.000 an. Das mediane Erkrankungsalter liegt bei ca. 75 Jahren, Frauen sind etwas seltener betroffen als Männer.

Die traditionell den MDS zugeordneten Typen werden in der aktuellen WHO-Klassifikation in 2 große Gruppen eingeteilt, siehe Tabelle 1. Neben den reinen MDS wird eine Gruppe von gemischten myelodysplastisch-myeloproliferativen Neoplasien abgegrenzt. Der von der akuten Leukämie diskriminierende Blasenanteil liegt in Blut und Knochenmark bei 20%. Der bis 2022 alleinig anwendbare Prognosescore für die MDS (IPSS-R) umfasst jedoch weiterhin Patienten mit bis zu 30% Blasten [4]. Für eine klare Diagnosestellung und Therapieentscheidung beim MDS ist eine Chromosomenanalyse und ein Screening auf somatische Mutationen unerlässlich, weil nur damit die Prognose des Patienten so gut als möglich bestimmt werden kann.

Den Vorschlägen der WHO-Klassifikation aus dem Jahre 2022 liegt ein neues Prinzip zugrunde, nämlich die Einteilung der MDS in morphologisch definierte und genetisch definierte Typen [1]. Zudem wurden drei neue MDS-Typen erstmals als eigenständige Entitäten definiert. Andere Typen wurden unverändert aus den alten WHO-Klassifikationen übernommen. Periphere Zellzahlen haben nun weniger Gewicht in der Klassifikation.

*Tabelle 3: Klassifikation der MDS nach den Vorschlägen der WHO 2022 [1]*

	Blastenanteil	Zytogenetik	Mutationen
<b>Genetisch definierte MDS</b>			
MDS mit niedrigen Blasten und isolierter Deletion (5q)	<5 % KM <2 % Blut	Deletion (5q) isoliert oder mit 1 anderen Anomalie außer Monosomie 7 oder Deletion (7q)	<i>SF3B1, TP53 möglich</i> <b>2 Subtypen</b> a) MDS del(5q) mit mono-allelischer <i>TP53</i> Mutation b) MDS del(5q) mit <i>SF3B1</i> Mutation
MDS mit niedrigen Blasten und <i>SF3B1</i> Mutation <sup>1</sup>	<5 % KM <2 % Blut	Keine Deletion (5q), keine Monosomie 7, kein komplex aberranter Karyotyp	Fast immer <i>SF3B1</i>
MDS mit bi-allelischer <i>TP53</i> Inaktivierung	Jeglicher	Typischerweise hoch komplex aberrant mit >3 Aberrationen	2 oder mehr <i>TP53</i> Mutationen oder 1 Mutation + Verlust der Kopienzahl von <i>TP53</i> <sup>5</sup>
<b>Morphologisch definierte MDS<sup>2</sup></b>			
<u>MDS mit niedrigen Blasten</u>			
MDS mit niedrigen Blasten <sup>3</sup>	<5% KM <2% Blut		
MDS, hypoplastisch <sup>4</sup>	<5% KM <2% Blut		

<u>MDS mit erhöhten Blasten</u>			
MDS mit erhöhten Blasten -1	5-9% KM, und/oder 2-4% Blut		
MDS mit erhöhten Blasten -2	10-19% KM und/oder 5-19% Blut		
MDS mit Fibrose	5-19% KM, und/oder 5-19% Blut		

<sup>1</sup> Nachweis von  $\geq 15\%$  Ringsideroblasten kann den Nachweis einer SF3B1-Mutation ersetzen

<sup>2</sup>  $\geq 10\%$  Dysplasiezeichen in mind. einer Zelllinie

<sup>3</sup> 2 Typen: MDS mit niedrigen Blasten und single lineage dysplasia (MDS-0-SLD); MDS mit niedrigen Blasten und multi lineage dysplasia (MDS-0-MLD)

<sup>4</sup>  $\leq 25\%$  histologische Knochenmarkzellularität, altersadaptiert

<sup>5</sup> Kopienzahlverlust entweder zytogenetisch (durch Bänderungsanalyse oder FISH darstellbare 17p-Deletion) oder Kopienzahl-neutraler Verlust der Heterozygotie (darstellbar mittels Array-Analysen oder spez. NGS-Verfahren). Bei einer VAF von 50% oder höher besteht ebenfalls eine hohe Wahrscheinlichkeit für eine biallelische TP53-Inaktivierung.

Bei Niedrigrisiko-MDS ist bei vielen MDS-Pat. aufgrund der geringgradigen Zytopenie zunächst eine „watch and wait“-Strategie ausreichend. Bei einem wesentlichen Teil der Fälle stellt jedoch die Anämie die häufigste Indikation zum Therapiebeginn dar. Eine Anämie führt vor allem bei älteren Pat. zu Fatigue, zu erhöhter Sturzhäufigkeit mit Frakturgefahr, zu verminderter Kognition und Lebensqualität sowie zu einem verkürzten Überleben.


Die Therapie mit Erythropoese stimulierenden Faktoren (ESF, klassisch: subkutanes Erythropoetin 40.000 IE/Woche, bei unzureichender Wirkung ggf. steigern auf 80.000 IE/Woche, einmal pro Woche; Verzögerungserthropoetin: 300 µg wöchentlich bzw. 500 µg zweiwöchentlich subkutan) muss in Anlehnung an den sogenannten „Nordic Score“ [4] erfolgen, siehe Abbildung 1. Die Kombination mit niedrigen Dosen von G-CSF (100 µg G-CSF s. c. 1-2 mal pro Woche mit dem Hintergrund, die Wirksamkeit von ESF zu modulieren, nicht um die Leukozyten anzuheben – s. o.) kann die Wirkung von ESF, insbesondere bei Pat. mit nachgewiesenen Ringsideroblasten, die refraktär auf eine alleinige ESF-Behandlung sind, verbessern.

Unter Berücksichtigung der prädiktiven Faktoren

- Erythropoetinspiegel <200 (500) U/l,
- geringe Transfusionsabhängigkeit (maximal 2 EK in 8 Wochen),
- niedriges Krankheitsrisiko,

kann ein Ansprechen bei bis zu 75 % der entsprechend ausgewählten Pat. erreicht werden [5, 6], siehe Abbildung 1. In der Regel ist das Ansprechen nach spätestens 6 Monaten Therapie zu erwarten. Bleibt es aus, sollte die Behandlung beendet werden. Ein Ansprechen ist bei einem Epo-Spiegel bis zu 500 U/l möglich.

Abbildung 1: Modifizierter Score der Nordic MDS-Group



Die Inhibition von (bisher teilweise nur unzureichend charakterisierten) Suppressoren der Erythropoese bei Patienten mit MDS führt insbesondere bei der Subgruppe der Pat. mit MDS und RS und/oder SF3B1-Mutation zu einer Verbesserung der Differenzierung und Steigerung der Proliferation von erythrozytären Zellen und damit zur Verringerung des Transfusionsbedarfes. Luspatercept, ein Inhibitor des TGF-beta Signalweges, ist in der Lage, bei etwa 60 % dieser transfusionsabhängigen Pat. eine signifikante Reduktion der Transfusionsbedürftigkeit bis hin zu Transfusionsfreiheit zu erzielen. Pat. mit MDS-RS (<5 % KM-Blasten, ≥15 % Ringsideroblasten im KM bzw. ≥5 % Ringsideroblasten im KM und Mutation von SF3B1) und einer transfusionsbedürftigen Anämie sollten mit Luspatercept behandelt werden, wenn sie auf ESF nicht angesprochen haben oder keine hohe Wahrscheinlichkeit des Ansprechens aufweisen (Serum-Epo-Spiegel ≥200 U/l) [7]. In dieser Zulassungsstudie wurde die Überlegenheit von Luspatercept gegenüber Placebo bei erwachsenen Patienten mit transfusionsabhängiger Anämie und MDS mit Ringsideroblasten mit sehr niedrigem, niedrigem oder intermediärem Risiko, die auf eine Erythropoetin-basierte Therapie nicht zufriedenstellend angesprochen haben oder dafür nicht geeignet sind, gezeigt. Die WHO 2022 Klassifikation führt formal das MDS mit Ringsideroblasten (MDS-RS) nicht mehr als eigene Entität auf, sondern hat die neue Entität MDS mit niedrigen Blasten und SF3B1 Mutation (siehe Tabelle) eingeführt wegen der prognostischen (günstigen) Rolle der o.g. Mutation. Die große Mehrzahl der Patienten mit MDS-RS weist eine solche Mutation auf, jedoch können auch Patienten mit MDS-RS ohne diese Mutation auf Luspatercept ansprechen [8].

Gibt es Kriterien für unterschiedliche Behandlungsentscheidungen in der o.g. Indikation, die regelhaft berücksichtigt werden? Wenn ja, welche sind dies und was sind in dem Fall die Therapieoptionen?  
(Bitte begründen Sie Ihre Ausführungen; geben Sie ggf. zitierte Quellen in einer Referenzliste an.)

Diese Kriterien sind oben aufgeführt.

#### Referenzliste:

- Khoury JD, Solary E, Abla O et al.: The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. Leukemia 36, 1703-1719, 2022. [DOI: 10.1038/s41375-022-01613-1](https://doi.org/10.1038/s41375-022-01613-1)

2. Hofmann WK et al.: Myelodysplastische Syndrome. ONKOPEDIA Gemeinsame Leitlinien von DGHO, OeGHO und SGH+SSH, Oktober 2020.  
<https://www.onkopedia.com/de/onkopedia/guidelines/myelodysplastische-syndrome-mds/@@guideline/html/index.html>
3. Neukirchen J, Schoonen WM, Strupp C et al.: Incidence and prevalence of myelodysplastic syndromes: data from the Düsseldorf MDS-registry. Leuk Res 35:1591-1596, 2011.  
<DOI:10.1016/j.leukres.2012.04.006>
4. Haferlach T, Nagata Y, Grossmann V et al.: Landscape of genetic lesions in 944 patients with myelodysplastic syndromes. Leukemia 28:241-247, 2014. <DOI:10.1038/leu.2013.336>
5. Hellström-Lindberg E, Negrin R, Stein R et al.: Erythroid response to treatment with G-CSF plus erythropoietin for the anaemia of patients with myelodysplastic syndromes: proposal for a predictive model. Br J Haematol 99:344-351, 1997. <DOI:10.1046/j.1365-2141.1997.4013211.x>
6. Platzbecker U, Symeonidis A, Olivia EN et al.: A phase 3 randomized placebo-controlled trial of darbepoetin alfa in patients with anemia and lower-risk myelodysplastic syndromes. Leukemia 31:1944-1950, 2017. <DOI:10.1038/leu.2017.192>
7. Fenaux P, Platzbecker U, Mufti GJ et al.: Luspatercept in patients with lower-risk myelodysplastic syndromes. N Engl J Med 382:140-151, 2020. <DOI:10.1056/NEJMoa1908892>
8. Platzbecker U, Haase D, Götze KS et al.: Long-Term Efficacy and Safety of Luspatercept for Anemia Treatment in Patients With Lower-Risk Myelodysplastic Syndromes: The Phase II PACE-MDS Study. J Clin Oncol 40:3800-3807, 2022. DOI: <10.1200/JCO.21.02476>